

**CCE RF
CCE RR**

A

ಕರ್ನಾಟಕ ಪ್ರೌಢ ಶಿಕ್ಷಣ ಪರೀಕ್ಷಾ ಮಂಡಳಿ, ಮಲ್ಲೇಶ್ವರಂ, ಬೆಂಗಳೂರು – 560 003

**KARNATAKA SECONDARY EDUCATION EXAMINATION BOARD, MALLESHWARAM,
BANGALORE – 560 003**

ಎಸ್.ಎಸ್.ಎಲ್.ಸಿ. ಪರೀಕ್ಷೆ, ಮಾರ್ಚ್ / ಏಪ್ರಿಲ್ — 2022

S. S. L. C. EXAMINATION, MARCH/APRIL, 2022

ಮಾದರಿ ಉತ್ತರಗಳು

MODEL ANSWERS

ದಿನಾಂಕ : 04. 04. 2022]

ಸಂಕೇತ ಸಂಖ್ಯೆ : **81-E**

Date : 04. 04. 2022]

CODE No. : **81-E**

ವಿಷಯ : ಗಣಿತ

Subject : MATHEMATICS

(ಶಾಲಾ ಅಭ್ಯರ್ಥಿ & ಪುನರಾವರ್ತಿತ ಶಾಲಾ ಅಭ್ಯರ್ಥಿ / Regular Fresh & Regular Repeater)

(ಇಂಗ್ಲಿಷ್ ಮಾಧ್ಯಮ / English Medium)

[ಗರಿಷ್ಠ ಅಂಕಗಳು : 80


[Max. Marks : 80

Qn. Nos.	Ans. Key	Value Points	Marks allotted
I. 1.		Multiple choice : $8 \times 1 = 8$ The graphical representation of the pair of lines $x + 2y - 4 = 0$ and $2x + 4y - 12 = 0$ is (A) intersecting lines (B) parallel lines (C) coincident lines (D) perpendicular lines. Ans. : (B) parallel lines	1


RF/RR (A)-(200)-9020 (MA)

[Turn over

Qn. Nos.	Ans. Key	Value Points	Marks allotted
2.		The common difference of the Arithmetic progression 8, 5, 2, - 1, ... is (A) - 3 (B) - 2 (C) 3 (D) 8. Ans. :	
	(A)	- 3	1
3.		The standard form of $2x^2 = x - 7$ is (A) $2x^2 - x = -7$ (B) $2x^2 + x - 7 = 0$ (C) $2x^2 - x + 7 = 0$ (D) $2x^2 + x + 7 = 0$. Ans. :	
	(C)	$2x^2 - x + 7 = 0$	1
4.		The value of $\cos (90^\circ - 30^\circ)$ is (A) - 1 (B) $\frac{1}{2}$ (C) 0 (D) 1. Ans. :	
	(B)	$\frac{1}{2}$	1
5.		The distance of the point $P(x, y)$ from the origin is (A) $\sqrt{x^2 + y^2}$ (B) $x^2 + y^2$ (C) $x^2 - y^2$ (D) $\sqrt{x^2 - y^2}$. Ans. :	
	(A)	$\sqrt{x^2 + y^2}$	1

Qn. Nos.	Ans. Key	Value Points	Marks allotted
6.		In a circle, the angle between the tangent and the radius at the point of contact is (A) 30° (B) 60° (C) 90° (D) 180°. Ans. :	
7.	(C)	90° In the given figure, the volume of the frustum of a cone is  (A) $\pi (r_1 + r_2) l$ (B) $\pi (r_1 - r_2) l$ (C) $\frac{1}{3} \pi h (r_1^2 - r_2^2 + r_1 r_2)$ (D) $\frac{1}{3} \pi h (r_1^2 + r_2^2 + r_1 r_2)$ Ans. :	1
8.	(D)	Surface area of a sphere of radius 'r' unit is (A) πr^2 sq.units (B) $2 \pi r^2$ sq.units (C) $3 \pi r^2$ sq.units (D) $4 \pi r^2$ sq.units. Ans. :	1
	(D)	$4 \pi r^2$ sq.units	1


Qn. Nos.	Value Points	Marks allotted
II.	Answer the following questions : $8 \times 1 = 8$ (Direct answers from Q. Nos. 9 to 16 full marks should be given)	
9.	If the pair of linear equations in two variables are inconsistent, then how many solutions do they have ? Ans. : No solution	1
10.	In an Arithmetic progression if 'a' is the first term and 'd' is the common difference, then write its n^{th} term. Ans. : $a_n = a + (n - 1)d$	1
11.	Write the standard form of quadratic equation. Ans. : $ax^2 + bx + c = 0$	1
12.	Write the value of $\frac{\sin 18^\circ}{\cos 72^\circ}$. Ans. : 1	1
13.	Write the distance of the point (4, 3) from x-axis. Ans. : 3	1
14.	Find the median of the scores 6, 4, 2, 10 and 7. Ans. : 6	1


Qn. Nos.	Value Points	Marks allotted
15.	Write the statement of “Basic Proportionality” theorem (Thales theorem). <i>Ans. :</i> If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, the other two sides are divided in the same ratio. Note : If correct alternate statement is written, give full marks.	1
16.	In the given figure, write the formula used to find the curved surface area of the cone.  <i>Ans. :</i> Curved surface area of cone = $\pi r l$ sq units	1
III.	Answer the following questions :	$8 \times 2 = 16$
17.	Solve the given pair of linear equations by Elimination method : $2x + y = 8$ $x - y = 1$ <i>Ans. :</i> $2x + y = 8 \dots\dots\dots (1)$ Adding $x - y = 1 \dots\dots\dots (2)$ $3x = 9$	$\frac{1}{2}$

Qn. Nos.	Value Points	Marks allotted
	$S_{20} = \frac{20}{2} [2(10) + (20-1)5]$ $= 10 [20 + 19 \times 5]$ $= 10 [20 + 95]$ $= 10 \times 115$ $S_{20} = 1150$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
	Note : Any other suitable method is followed to get the correct answer, full marks should be given. OR $S_n = \frac{n(n+1)}{2}$ $n = 20$ $S_{20} = \frac{20(20+1)}{2}$ $= \frac{20 \times 21}{2}$ $= 10 \times 21$ $S_{20} = 210$	 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 2
20.	Find the roots of $x^2 + 5x + 2 = 0$ by using quadratic formula. Ans. : $x^2 + 5x + 2 = 0$ $ax^2 + bx + c = 0$ $a = 1, b = 5, c = 2$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $= \frac{-5 \pm \sqrt{5^2 - 4(1)(2)}}{2(1)}$ $= \frac{-5 \pm \sqrt{25 - 8}}{2}$ $= \frac{-5 \pm \sqrt{17}}{2}$	 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 2

Qn. Nos.	Value Points	Marks allotted
21.	Find the value of the discriminant and hence write the nature of roots of the quadratic equation $x^2 + 4x + 4 = 0$. Ans. : $x^2 + 4x + 4 = 0$ $ax^2 + bx + c = 0$ $a = 1, b = 4, c = 4$ $\text{Discriminant} = b^2 - 4ac$ $= 4^2 - 4(1)(4)$ $= 16 - 16$ $= 0$ Nature of roots : Two equal real roots.	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 2
22.	Find the distance between the points A (2, 6) and B (5, 10) by using distance formula. OR Find the coordinates of the mid-point of the line segment joining the points P (3, 4) and Q (5, 6) by using 'mid-point' formula. Ans. : A (2, 6) B (5, 10) $x_1, y_1 \quad x_2, y_2$ $\text{Distance formula } d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ $= \sqrt{(5 - 2)^2 + (10 - 6)^2}$ $= \sqrt{3^2 + 4^2}$ $= \sqrt{9 + 16}$ $= \sqrt{25}$ $d = 5 \text{ units}$ OR P (3, 4) Q (5, 6) $x_1, y_1 \quad x_2, y_2$ $\text{Mid-point formula } P (x, y) = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 2 $\frac{1}{2}$

Qn. Nos.	Value Points	Marks allotted
	Ans. : (i) $\sin \theta = \frac{12}{13}$ 1 (ii) $\tan \alpha = \frac{5}{12}$ 1	2
IV.	Answer the following questions : $9 \times 3 = 27$	
25.	The sum of first 9 terms of an Arithmetic progression is 144 and its 9th term is 28 then find the first term and common difference of the Arithmetic progression. Ans. : $S_n = \frac{n}{2} [a + l]$ $\frac{1}{2}$ $S_9 = \frac{9}{2} [a + 28]$ $144 = \frac{9}{2} [a + 28]$ $\frac{1}{2}$ $\frac{144 \times 2}{9} = a + 28$ $\left. \begin{array}{l} \frac{16}{9} \times 2 \\ 144 \times 2 \\ 9 \end{array} \right\}$ $32 = a + 28$ $a = 32 - 28$ $\left. \begin{array}{l} a = 32 - 28 \\ a = 4 \end{array} \right\}$ $\frac{1}{2}$ $a_n = a + (n - 1) d$ $\frac{1}{2}$ $a_9 = 4 + (9 - 1) d$ $\left. \begin{array}{l} a_9 = 4 + (9 - 1) d \\ 28 = 4 + 8d \end{array} \right\}$ $\frac{1}{2}$ $28 = 4 + 8d$ $24 = 8d$ $d = \frac{24}{8}$ $d = 3$ $\frac{1}{2}$ * Any other correct alternate, method may be given full marks.	3

Qn. Nos.	Value Points	Marks allotted
26.	The diagonal of a rectangular field is 60 m more than its shorter side. If the longer side is 30 m more than the shorter side, then find the sides of the field. OR In a right angled triangle, the length of the hypotenuse is 13 cm. Among the remaining two sides, the length of one side is 7 cm more than the other side. Find the sides of the triangle. Ans. :  $ABCD \rightarrow$ rectangular field Let $AB = x$ m then $BC = (x + 30)$ m, $AC = (x + 60)$ m $AC^2 = AB^2 + BC^2$ $(x + 60)^2 = x^2 + (x + 30)^2$ $\cancel{x^2} + 60^2 + 2 \times x \times 60 = \cancel{x^2} + x^2 + 30^2 + 2 \times x \times 30$ $3600 + 120x = x^2 + 900 + 60x$ $x^2 + 900 + 60x - 3600 - 120x = 0$ $x^2 - 60x - 2700 = 0$ $x^2 - 90x + 30x - 2700 = 0$ $x(x - 90) + 30(x - 90) = 0$ $(x - 90)(x + 30) = 0$ $x - 90 = 0 \quad \text{or} \quad x + 30 = 0$ $x = 90 \quad \text{or} \quad x = -30 \quad (\text{not considered})$ $\therefore x = 90$ $AB = x = 90 \text{ m}$ $BC = (x + 30) = 90 + 30 = 120 \text{ m}$ OR	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 3

Qn. Nos.	Value Points	Marks allotted
	 Let ABC be a right angled triangle. Let $AC = 13$ cm, $AB = x$ cm and $BC = (x + 7)$ cm $AC^2 = AB^2 + BC^2$ $13^2 = x^2 + (x + 7)^2$ $\Rightarrow 169 = x^2 + x^2 + 49 + 14x$ $\Rightarrow 169 = 2x^2 + 49 + 14x$ $\Rightarrow 2x^2 + 49 + 14x - 169 = 0$ $\Rightarrow 2x^2 + 14x - 120 = 0$ $\div 2, \quad x^2 + 7x - 60 = 0$ $\Rightarrow x^2 + 12x - 5x - 60 = 0$ $\Rightarrow x(x + 12) - 5(x + 12) = 0$ $\Rightarrow (x + 12)(x - 5) = 0$ $x + 12 = 0 \text{ or } x - 5 = 0$ $x = -12 \text{ or } x = 5$ (not considered) $\therefore x = 5$ $AB = x = 5 \text{ cm}$ $BC = (x + 7) = 5 + 7 = 12 \text{ cm}$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 3


Qn. Nos.	Value Points	Marks allotted
27.	Prove that $(\sin A + \operatorname{cosec} A)^2 + (\cos A + \sec A)^2 = 7 + \tan^2 A + \cot^2 A.$ OR Prove that : $\sec \theta (1 - \sin \theta) (\sec \theta + \tan \theta) = 1.$ Ans. : LHS = $(\sin A + \operatorname{cosec} A)^2 + (\cos A + \sec A)^2$ $= \sin^2 A + \operatorname{cosec}^2 A + 2 \sin A \cdot \operatorname{cosec} A + \cos^2 A + \sec^2 A + 2 \cos A \cdot \sec A$ $= \sin^2 A + \cos^2 A + \operatorname{cosec}^2 A + 2 \cancel{\sin A} \cdot \frac{1}{\cancel{\sin A}} + \sec^2 A + 2 \cancel{\cos A} \cdot \frac{1}{\cancel{\cos A}}$ $= 1 + (1 + \cot^2 A) + 2 + (1 + \tan^2 A) + 2$ $[\because \operatorname{cosec}^2 A = 1 + \cot^2 A$ $\sec^2 A = 1 + \tan^2 A$ $\sin^2 A + \cos^2 A = 1]$ $= 7 + \tan^2 A + \cot^2 A$ LHS = RHS OR LHS = $\sec \theta (1 - \sin \theta) (\sec \theta + \tan \theta)$ $= \frac{1}{\cos \theta} (1 - \sin \theta) \left(\frac{1}{\cos \theta} + \frac{\sin \theta}{\cos \theta} \right)$ $= \frac{(1 - \sin \theta)}{\cos \theta} \times \frac{(1 + \sin \theta)}{\cos \theta}$ $= \frac{1 - \sin^2 \theta}{\cos^2 \theta}$ $= \frac{\cos^2 \theta}{\cos^2 \theta} [\because 1 - \sin^2 \theta = \cos^2 \theta]$ $= 1$ $\therefore$ L.H.S. = R.H.S	1 1 1/2 1/2 3 1 1/2 1/2 1/2 1/2 3


Qn. Nos.	Value Points	Marks allotted
28.	Find the coordinates of the point on the line segment joining the points $A (-1, 7)$ and $B (4, -3)$ which divides AB internally in the ratio $2 : 3$. OR Find the area of triangle PQR with vertices $P (0, 4)$, $Q (3, 0)$ and $R (3, 5)$. Ans. : $A (-1, 7),$ x_1, y_1 $B (4, -3)$ x_2, y_2 $2 : 3$ $m_1 \ m_2$ $P(x, y) = \left(\frac{m_1 x_2 + m_2 x_1}{m_1 + m_2}, \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} \right)$ $= \left(\frac{2(4) + 3(-1)}{2+3}, \frac{2(-3) + 3(7)}{2+3} \right)$ $= \left(\frac{8-3}{5}, \frac{-6+21}{5} \right)$ $= \left(\frac{5}{5}, \frac{15}{5} \right)$ $P(x, y) = (1, 3)$ OR $P (0, 4),$ x_1, y_1 $Q (3, 0)$ x_2, y_2 $R (3, 5)$ x_3, y_3 $A = \frac{1}{2} [x_1 (y_2 - y_3) + x_2 (y_3 - y_1) + x_3 (y_1 - y_2)]$ $= \frac{1}{2} [0(0-5) + 3(5-4) + 3(4-0)]$	1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 3

Qn. Nos.	Value Points	Marks allotted												
	$= \frac{1}{2}[0(-5)+3(1)+3(4)]$	$\frac{1}{2}$												
	$= \frac{1}{2} [0 + 3 + 12]$	$\frac{1}{2}$												
	$= \frac{1}{2} \times 15$													
	$A = \frac{15}{2} \text{ or } 7.5 \text{ sq. units}$	$\frac{1}{2}$												
	29. Find the mean for the following grouped data by Direct method :													
	<table><tr><th><i>Class-interval</i></th><th><i>Frequency</i></th></tr><tr><td>10 — 20</td><td>2</td></tr><tr><td>20 — 30</td><td>3</td></tr><tr><td>30 — 40</td><td>5</td></tr><tr><td>40 — 50</td><td>7</td></tr><tr><td>50 — 60</td><td>3</td></tr></table>	<i>Class-interval</i>	<i>Frequency</i>	10 — 20	2	20 — 30	3	30 — 40	5	40 — 50	7	50 — 60	3	
<i>Class-interval</i>	<i>Frequency</i>													
10 — 20	2													
20 — 30	3													
30 — 40	5													
40 — 50	7													
50 — 60	3													
	OR													
	Find the mode for the following grouped data :													
	<table><tr><th><i>Class-interval</i></th><th><i>Frequency</i></th></tr><tr><td>5 — 15</td><td>3</td></tr><tr><td>15 — 25</td><td>4</td></tr><tr><td>25 — 35</td><td>8</td></tr><tr><td>35 — 45</td><td>7</td></tr><tr><td>45 — 55</td><td>3</td></tr></table>	<i>Class-interval</i>	<i>Frequency</i>	5 — 15	3	15 — 25	4	25 — 35	8	35 — 45	7	45 — 55	3	
<i>Class-interval</i>	<i>Frequency</i>													
5 — 15	3													
15 — 25	4													
25 — 35	8													
35 — 45	7													
45 — 55	3													

Qn. Nos.	Value Points	Marks allotted																												
	Ans. : <table><tr><th>C-I</th><th>f_i</th><th>x_i</th><th>$f_i x_i$</th></tr><tr><td>10-20</td><td>2</td><td>15</td><td>30</td></tr><tr><td>20-30</td><td>3</td><td>25</td><td>75</td></tr><tr><td>30-40</td><td>5</td><td>35</td><td>175</td></tr><tr><td>40-50</td><td>7</td><td>45</td><td>315</td></tr><tr><td>50-60</td><td>3</td><td>55</td><td>165</td></tr><tr><td></td><td>$N = 20$</td><td></td><td>$\sum f_i x_i = 760$</td></tr></table> Table 2 [Mid points – 01 finding $f_i x_i$ - 01] Mean, $\bar{X} = \frac{\sum f_i x_i}{N}$ OR $\frac{\sum FX}{N}$ 1/2 $= \frac{760}{20}$ $\bar{X} = 38$ 1/2 OR From the frequency distribution table we find that $f_0 = 4, f_1 = 8, f_2 = 7, h = 10$ and $l = 25$ Mode = $l + \left[\frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right] \times h$ 1 $= 25 + \left[\frac{8 - 4}{2(8) - 4 - 7} \right] \times 10$ 1/2 $= 25 + \left[\frac{4}{16 - 11} \right] \times 10$ 1/2 $= 25 + \frac{4}{5} \times 10^2$ 1/2 $= 25 + 8$ Mode = 33 1/2	C-I	f_i	x_i	$f_i x_i$	10-20	2	15	30	20-30	3	25	75	30-40	5	35	175	40-50	7	45	315	50-60	3	55	165		$N = 20$		$\sum f_i x_i = 760$	3
C-I	f_i	x_i	$f_i x_i$																											
10-20	2	15	30																											
20-30	3	25	75																											
30-40	5	35	175																											
40-50	7	45	315																											
50-60	3	55	165																											
	$N = 20$		$\sum f_i x_i = 760$																											



Qn. Nos.	Value Points	Marks allotted														
30.	During a medical check-up of 50 students of a class, their heights were recorded as follows : Draw “less than type” ogive for the given data : <table><tr><th>Height in cm</th><th>Number of students (Cumulative frequency)</th></tr><tr><td>Less than 140</td><td>5</td></tr><tr><td>Less than 145</td><td>10</td></tr><tr><td>Less than 150</td><td>15</td></tr><tr><td>Less than 155</td><td>25</td></tr><tr><td>Less than 160</td><td>40</td></tr><tr><td>Less than 165</td><td>50</td></tr></table> Ans. : Scale x-axis 1 cm = 5 units y-axis 1 cm = 5 units The graph shows a smooth curve (ogive) plotted on a coordinate system. The x-axis is labeled 'upper limits' and has tick marks at 140, 145, 150, 155, 160, and 165. The y-axis is labeled 'cumulative frequency' and has tick marks from 0 to 50 in increments of 5. The curve starts at the point (140, 5) and ends at (165, 50), passing through (145, 10), (150, 15), (155, 25), and (160, 40). A scale is provided: x-axis 1 cm = 5 units, y-axis 1 cm = 5 units.	Height in cm	Number of students (Cumulative frequency)	Less than 140	5	Less than 145	10	Less than 150	15	Less than 155	25	Less than 160	40	Less than 165	50	
Height in cm	Number of students (Cumulative frequency)															
Less than 140	5															
Less than 145	10															
Less than 150	15															
Less than 155	25															
Less than 160	40															
Less than 165	50															

Qn. Nos.	Value Points	Marks allotted
	Drawing axes and writing scale $\frac{1}{2} + \frac{1}{2} = 1$	
	Marking points 1	
	Drawing Ogive 1	3
31.	Prove that “the lengths of tangents drawn from an external point to a circle are equal”. Ans. :  Data : O is the centre of the circle. PQ and PR are tangents drawn from external point 'P'. To Prove : $PQ = PR$ Construction : Join OP, OQ and OR. Proof : In the figure $\angle OQP = \angle ORP = 90^\circ \quad \left[\begin{array}{l} OQ \perp PQ \\ OR \perp PR \end{array} \right]$ $\begin{array}{ll} OQ = OR & [\text{radii of same circle}] \\ OP = OP & [\text{common side}] \\ \Delta OQP \cong \Delta ORP & [\text{RHS}] \\ PQ = PR & [\text{CPCT}] \end{array} \quad \left. \vphantom{\begin{array}{l} OQ = OR \\ OP = OP \\ \Delta OQP \cong \Delta ORP \\ PQ = PR \end{array}} \right\}$ Note : If the theorem is proved as given in the text-book, give full marks.	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 3



Qn. Nos.	Value Points	Marks allotted
32.	Construct two tangents to a circle of radius 3 cm from a point 8 cm away from its centre. Ans. :  Drawing a circle C_1 of radius 3 cm $\frac{1}{2}$ Drawing $OP = 8$ cm $\frac{1}{2}$ Constructing perpendicular bisector of OP 1 Drawing C_2 circle $\frac{1}{2}$ Joining PA and PB $\frac{1}{2}$	3
33.	The volume of a solid right circular cylinder is 2156 cm^3. If the height of the cylinder is 14 cm, then find its curved surface area. [Take $\pi = \frac{22}{7}$]	

Qn. Nos.	Value Points	Marks allotted
	Ans. : $V = 2156 \text{ cm}^3$ $h = 14 \text{ cm}$ $r = ?$ CSA = ? Volume of cylinder = $\pi r^2 h$ $\frac{1}{2}$ $2156 = \frac{22}{7} \times r^2 \times 14$ $\frac{1}{2}$ $2156 = 44 r^2$ $r^2 = \frac{2156}{44}$ $r^2 = 49$ $r = \sqrt{49}$ $r = 7 \text{ cm}$ $\frac{1}{2}$ Curved surface area of cylinder $\left\{ \begin{array}{l} = 2\pi rh \\ = 2 \times \frac{22}{7} \times 7 \times 14 \end{array} \right.$ $\frac{1}{2}$ $= 2 \times 22 \times 14$ $\frac{1}{2}$ $= 616 \text{ cm}^2$ $\frac{1}{2}$	3
V.	Answer the following questions :	$4 \times 4 = 16$
34.	Find the solution of the given pair of linear equations by graphical method : $x + 2y = 6$ $x + y = 5$	


Qn. Nos.	Value Points	Marks allotted												
Ans. :	Scale x-axis 1 cm = 1 unit y-axis 1 cm = 1 unit $(4, 1)$ $x + 2y = 6$ $x + y = 5$ $x = 4$ $y = 1$ $x + 2y = 6$ <table border="1"> <tr><td>x</td><td>0</td><td>6</td></tr> <tr><td>y</td><td>3</td><td>0</td></tr> </table> $x + y = 5$ <table border="1"> <tr><td>x</td><td>0</td><td>5</td></tr> <tr><td>y</td><td>5</td><td>0</td></tr> </table> For table construction (1 + 1) 2 Drawing two lines by marking points 1 Marking point of intersection & writing values of x and y 1 Note : Any other points also may be considered to get straight lines. 4	x	0	6	y	3	0	x	0	5	y	5	0	
x	0	6												
y	3	0												
x	0	5												
y	5	0												
35.	The angle of elevation of the top of a building from the foot of a tower is 30° and the angle of elevation of the top of the tower from the foot of the building is 60°. Both the tower and building are on the same level													

Qn. Nos.	Value Points	Marks allotted
	ground. If the height of the tower is 50 m, then find the height of the building.  OR As observed from the top of a 75 m high light house from the sea-level, the angles of depression of two ships are 30° and 45°. If one ship is exactly behind the other on the same side of the light house, then find the distance between the two ships.  Ans. : In $\triangle BDC$, $\tan 60^\circ = \frac{CD}{BD}$ $\frac{1}{2}$ $\sqrt{3} = \frac{50}{BD}$ $\therefore BD = \frac{50}{\sqrt{3}} \dots\dots\dots (1)$ In $\triangle ABD$, $\tan 30^\circ = \frac{AB}{BD}$ $\frac{1}{2}$	

Qn. Nos.	Value Points	Marks allotted
	$\frac{1}{\sqrt{3}} = \frac{AB}{BD}$ $BD = \sqrt{3} \cdot AB \dots\dots\dots (2)$ $\frac{1}{2}$ From (1) and (2) $\sqrt{3} \cdot AB = \frac{50}{\sqrt{3}}$ $\frac{1}{2}$ $AB = \frac{50}{\sqrt{3} \cdot \sqrt{3}}$ $\frac{1}{2}$ $AB = \frac{50}{3} \text{ or } 16\frac{2}{3} \text{ m}$ $\frac{1}{2}$ OR Distance between the two ships is PQ In $\triangle ABP$, $\tan 45^\circ = \frac{AB}{BP}$ $\frac{1}{2}$ $1 = \frac{75}{BP}$ $\frac{1}{2}$ $\therefore BP = 75$ $\frac{1}{2}$ In $\triangle ABQ$, $\tan 30^\circ = \frac{AB}{BQ}$ $\frac{1}{2}$ $\frac{1}{\sqrt{3}} = \frac{75}{BP + PQ}$ $\frac{1}{2}$ $\frac{1}{\sqrt{3}} = \frac{75}{75 + PQ}$ $\frac{1}{2}$ $75 + PQ = 75\sqrt{3}$ $PQ = 75\sqrt{3} - 75$ $\frac{1}{2}$ $PQ = 75(\sqrt{3} - 1) \text{ m}$ $\frac{1}{2}$	4
36.	Construct a triangle with sides 4.5 cm, 6 cm and 8 cm. Then construct another triangle whose sides are $\frac{3}{4}$ of the corresponding sides of the first triangle.	4

Qn. Nos.	Value Points	Marks allotted
	Ans. :  Construction of given triangle 1 Construction of acute angle with division 1 Drawing parallel lines 1 Obtaining required triangle 1	4
37.	In the figure AXB and CYD are the arcs of two concentric circles with centre O. The length of the arc AXB is 11 cm. If $OC = 7$ cm and $\angle AOB = 30^\circ$, then find the area of the shaded region. [Take $\pi = \frac{22}{7}$] 	

Qn. Nos.	Value Points	Marks allotted
	Ans. : Length of the arc = $\frac{\theta}{360^\circ} \times 2\pi r$ $\frac{1}{2}$ $11 = \frac{30^\circ}{360^\circ} \times 2 \times \frac{22^{11}}{7} \times r$ $\frac{1}{2}$ $11 = \frac{11r}{21}$ $r = \frac{11 \times 21}{11}$ $r = 21 \text{ cm}$ $\frac{1}{2}$ Area of the sector OAXB = $A_1 = \frac{\theta}{360^\circ} \times \pi r^2$ $\frac{1}{2}$ $= \frac{30^\circ}{360^\circ} \times \frac{22}{7} \times 21^2$ $= \frac{1}{12} \times \frac{22^{11}}{7} \times 21^1 \times 21$ $= \frac{231}{2} \text{ cm}^2$ $\frac{1}{2}$ Area of the sector OCYD = $A_2 = \frac{\theta}{360^\circ} \times \pi r^2$ $= \frac{30^\circ}{360^\circ} \times \frac{22}{7} \times 7^2$ $= \frac{1}{12} \times \frac{22^{11}}{7} \times 7 \times 7$ $A_2 = \frac{77}{6} \text{ cm}^2$ $\frac{1}{2}$ Area of the shaded region = $A_1 - A_2$ $= \frac{231}{2} - \frac{77}{6}$ $= \frac{693 - 77}{6}$ $\frac{1}{2}$ $= \frac{616}{6}$	

Qn. Nos.	Value Points	Marks allotted
	$= \frac{308}{3} \text{ cm}^2$ or $= 102.66 \text{ cm}^2$ $\text{OR } 102.7 \text{ cm}^2$	$\frac{1}{2}$ 4
VI.	Answer the following question :	$1 \times 5 = 5$
38.	Prove that “the ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding sides”. Ans. :  $\frac{1}{2}$ Data : $\triangle ABC \sim \triangle PQR$ $\therefore \frac{AB}{PQ} = \frac{BC}{QR} = \frac{AC}{PR}$ $\frac{1}{2}$ To prove : $\frac{\text{Area of } \triangle ABC}{\text{Area of } \triangle PQR} = \frac{BC^2}{QR^2}$ $\frac{1}{2}$ Construction : Draw $AM \perp BC$ and $PN \perp QR$ $\frac{1}{2}$ Proof : $\frac{\text{Area of } \triangle ABC}{\text{Area of } \triangle PQR} = \frac{\frac{1}{2} \times BC \times AM}{\frac{1}{2} \times QR \times PN}$ $\frac{1}{2}$ $\frac{\text{Area of } \triangle ABC}{\text{Area of } \triangle PQR} = \frac{BC}{QR} \times \frac{AM}{PN} \dots\dots\dots (1)$ $\frac{1}{2}$ In $\triangle ABM$ and $\triangle PQN$ $\angle B = \angle Q$	

RF/RR (A)-(200)-9020 (MA)