Board Of School Education Haryana



मॉडल पेपर

उत्तर

2025


aglasem.com





MARKING SCHEME BSEH PRACTICE PAPER 2, 10TH MATHS(BASIC) , March2025 (ENGLISH MEDIUM)		
Q. no.	**Expected solutions**	**marks**
	Section-A	
1	(d)60	1
2	(d)more than 3	1
3	(c)$(x+2)(x-1)=x^2-2x-3$	1
4	(c)3 units	1
5	(a) -12	1
6	(a) 50°	1
7	(d) 55°	1
8	(b) $\frac{b}{\sqrt{a^2+b^2}}$	1
9	(a)60^0	
10	(b) $10\sqrt{2}$	
11	(d) 3	1
12	(a) $\frac{1}{5}$	1
13	Irrational number	1
14	$\sqrt{119}$ cm	1
15	tanθ =a b	1
16	$\frac{1}{2}$	1
17	$\frac{77}{2}$ cm² or $\frac{49\pi}{4}$ cm²	1
18	False	1
19	(a)Both Assertion(A) and Reason (R) are true and Reason (R) is the correct explanation of Assertion(A).	1
20	(b) Both Assertion(A) and Reason (R) are true but Reason (R) is the not correct explanation of Assertion(A).	1
	SECTION-B	
21. (a)	x/2 + 2y/3 = -1 3x + 4y = -6 ..... (i)	1/2

	…… x-y/3 = 3 3x – y = 9 …… (ii) …… When the equation (ii) is subtracted from equation (i) we get, 5y = -15 y = -3 ……(iii) …… When the equation (iii) is substituted in (i) we get, 3x – 12 = -6 3x = 6 x = 2 Hence, x = 2 , y = -3	1/2 1/2 1/2
21. (b)	Using the property of a rectangle, We know that, Lengths are equal, i.e., CD = AB Hence, x + 3y = 13 …(i) …… Breadths are equal, i.e., AD = BC Hence, 3x + y = 7 …(ii) …… On multiplying Eq. (ii) by 3 and then subtracting Eq. (i), We get, 8x = 8 So, x = 1 …… On substituting x = 1 in Eq. (i), We get, y = 4 Therefore, the required values of x and y are 1 and 4, respectively.	1/2 1/2 1/2 1/2
22.	Let (– 4, 6)divide AB internally in the ratio k : 1. Using the section formula, we get $(-4, 6) = \left(\frac{3k-6}{k+1}, \frac{-8k+10}{k+1}\right)$ ……	1

	So, $-4 = \frac{3k-6}{k+1}$ …… $\Rightarrow -4k - 4 = 3k - 6$ $\Rightarrow 7k = 2$ $\Rightarrow$, k : 1 = 2 : 7 We can check for the y-coordinate also. So, the point (– 4, 6) divides the line segment joining the points A(– 6, 10) and B(3, – 8) in the ratio 2 : 7.	1/2 1/2
23.	 In ΔPBC and ΔPDE, ∠BPC = ∠EPD [vertically opposite angles] PB/PD = 5/10 = ½ … (i) PC/PE = 6/12 = ½ … (ii) …… From equation (i) and (ii), We get, PB/PD = PC/PE Since, ∠BPC of ΔPBC = ∠EPD of ΔPDE and the sides including these. …… Then, by SAS similarity criteria ΔPBC ~ ΔPDE	 1/2 1/2
24.	We know that,	

(b)	cos 60° = 1/2 sec 30° = 2/√3 tan 45° = 1 sin 30° = 1/2 cos 30° = √3/2 ……	1/2
	Now, substitute the values in the given problem, we get $(5\cos^2 60° + 4\sec^2 30° - \tan^2 45°)/(\sin^2 30° + \cos^2 30°)$ $= \{5(1/2)^2+4(2/\sqrt{3})^2-1\}/(1/2)^2+(\sqrt{3}/2)^2$ ……	1/2
	= (5/4+16/3-1)/(1/4+3/4) ={ (15+64-12)/12}/(4/4) ……	1
	= 67/12	1/2
24. (a)	LHS = $\sqrt{\frac{1+\sin A}{1-\sin A}} =$ $= \sqrt{\frac{1+sinA}{1-sinA} \times \frac{1+sinA}{1+sinA}}$ ……	1/2
	$= \frac{1+\sin A}{\sqrt{1-\sin^2 A}}$ ……	1/2

	$= \frac{1+\sin A}{\sqrt{\cos^2 A}}$ …… $= \frac{1+\sin A}{\cos A}$ = secA +tanA = RHS	1/2 1/2
25.	Area swept by the minute hand in 60 minutes = Area of the circle with radius equal to the length of the minute hand = πr^2 …… Area swept by minute hand in 1 minute = $\pi r^2/60$ …… Thus, area swept by minute hand in 5 minutes = $(\pi r^2/60) \times 5 = \pi r^2/12$ [∵Length of the minute hand (r) = 14 cm] …… = 1/12 × 22/7 × 14 × 14 cm^2 = 154/3 cm^2	1/2 1 1/2 1/2
	SECTION-C	


26.	Prove that $\sqrt{2}$ is irrational. Solution: Let, if possible,$\sqrt{2}$ be a rational no. --- $\therefore \sqrt{2} = \frac{p}{q}$, where p and q are co-prime integers and $q \neq 0$. --- $\Rightarrow 2 = \frac{p^2}{q^2}$ $\Rightarrow p^2 = 2q^2$ .....(i) $\Rightarrow$ 2 divides $p^2 \Rightarrow$ 2 divides p also. --- Let p = 2m,.....(ii) where m is any integer. $\Rightarrow p^2 = 4m^2$.....(iii) --- From (ii) and (iii) $2q^2 = 4m^2$ $\Rightarrow q^2 = 2m^2$ $\Rightarrow$ 2 divides $q^2 \Rightarrow$2 divides q also. $\Rightarrow$ q = 2n.....(iv) --- From (i) and (iv) , p and q have 2 as common factor. $\therefore$ p and q are not co-prime. Hence our supposition is wrong. $\therefore \sqrt{2}$ is an irrational number.	1/2 1/2 1/2 1 1/2 1/2
27.	$6x^2-3-7x=6x^2-7x-3=0$ $\Rightarrow 6x^2+2x-9x-3=0$	

	$\Rightarrow 2x(3x+1)-3(3x+1)=0$ $\Rightarrow (2x-3)(3x+1)=0$ Zeros = $3/2, -1/3$ ……	1
	$\alpha+\beta=-b/a \Rightarrow (3/2)+(-1/3)=7/6=-(-7)6=-b/a$ ……	1
	$\alpha\beta=c/a \Rightarrow (3/2)(-1/3)=-1/2=-3/6=c/a$ Hence proved.	1
28. (a)	Let Rahul's age be x years and his son's age be y years. …… Five years hence(later), $x+5=3(y+5)$ $\Rightarrow x+5=3y+15$ $\Rightarrow x-3y=10$.....(1) …… Also, five years ago(before), $x-5=7(y-5)$ $\Rightarrow x-5=7y-35$ $\Rightarrow x-7y=-30$.....(2) …… Subtracting equation (2) from (1), $x-3y-x+7y = 10+30$ (∵ eq.(2) changes its sign)	1/2  1/2 1/2 1/2


	$4y = 40$ $\Rightarrow y = 10$ …… Put y = 10 in eq. (1), $x - 3(10) = 10$ $\Rightarrow x - 30 = 10$ $\Rightarrow x = 40$ …… Thus, present age of Rahul=x=40 years and present age of Rahul's son=y=10 years.	1/2 1/2
28. (b)	Let the larger angle = x Smaller angle = y As both angles are supplementary, $x + y = 180$ $\Rightarrow x = 180 - y$ .... (i) …… Difference is 18 degrees. So, $x - y = 18$ $\Rightarrow x = 18 + y$ .... (i) …… Substituting the value of x in equation (i) we get, $\Rightarrow 18 + y = 180 - y$ $\Rightarrow - y - y = 18 - 180$ $\Rightarrow - 2y = -162$ $\Rightarrow y = -162/-2$ $\Rightarrow y = 81$ …… Substituting the value of y in equation (i), we get, $\Rightarrow x = 180 - 81 = 99$	 1/2 1/2 1/2

	…… Hence, the angles are 99° and 81°.	1/2
29.	We know that the distance between the two points is given by the Distance Formula = $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ By substituting the values of points P (2, - 3) and Q (10, y) in the distance formula, we get …… PQ = $\sqrt{(2 - 10)^2 + (-3 - y)^2} = 10$ PQ = $\sqrt{(-8)^2 + (3 + y)^2} = 10$ …… Squaring on both sides, we get $64 + (y + 3)^2 = 100$ …… $(y + 3)^2 = 36$ $y + 3 = \sqrt{36}$ $y + 3 = \pm 6$ …… y + 3 = 6 or y + 3 = - 6 Therefore, y = 3 or - 9 are the possible values for y.	1/2 1/2 1/2 1/2 1
30. (a)	Given, $\cos A + \cos^2 A = 1$ $\Rightarrow \cos A = 1 - \cos^2 A$ $\Rightarrow \cos A = \sin^2 A$ [∵ $\sin^2 A = 1 - \cos^2 A$] ……(i) …… LHS=$(\sin^2 A + \sin^4 A)$ = $(\sin^2 A + (\sin^2 A)^2)$ ……	1 1/2


	$= (\sin^2 A + (\cos A)^2)$ [using (i)] …… $= \sin^2 A + \cos^2 A$ $= 1$ = RHS	1 1/2
30. (b)	LHS=$(\sin A+\operatorname{cosec} A)^2+(\cos A+\sec A)^2$ $=\sin^2 A+\operatorname{cosec}^2 A+2\sin A\operatorname{cosec} A+\cos^2 A+\sec^2 A+2\cos A\sec A$ …… $=\sin^2 A+\cos^2 A+\operatorname{cosec}^2 A+\sec^2 A+2\sin A\times 1/\sin A+2\cos A\times 1/\cos A$ [∵cosecA=1/sinA and secA=1/cosA] …… $=1+\operatorname{cosec}^2 A+\sec^2 A+2+2$ [∵$\sin^2 A+\cos^2 A=1$] …… $=5+(1+\cot^2 A)+(1+\tan^2 A)$ [∵$1+\tan^2 A=\sec^2 A$ and $1+\cot^2 A=\operatorname{cosec}^2 A$] $=7+\tan^2 A+\cot^2 A$= RHS ……	1/2 1 1/2 1


31.	 ……. Given the height of the observer be DE = 1.5 m That is AB = 1.5 m Let BC = h is the height of the chimney Hence AC = (h – 1.5) m Given the distance between the observer and the chimney is AD = BE = 28.5 m ……. In right $\triangle CAD, \theta = 45^{o}$ $\tan 45^{o} = AC/AD$ $\Rightarrow 1 = (h-1.5)/28.5$ ……. $\Rightarrow 28.5 = h - 1.5$ $\Rightarrow h = 28.5 + 1.5 = 30$ m Thus the height of the chimney is 30 m.	1/2 1 1 1
	SECTION-D	
32. (a)	Given, 2nd term, $a_2 = 14$ 3rd term, $a_3 = 18$	1

	Common difference, $d = a_3 - a_2 = 18 - 14 = 4$ ..... We know that nth term of an AP is, $a_n = a + (n - 1)d$ $a_2 = a + d$ $14 = a + 4$ $a = 10$ ..... Sum of n terms of AP is given by $S_n = n/2 [2a + (n - 1) d]$ ..... $S_{51} = 51/2 [2 \times 10 + (51 - 1) 4]$ ..... $= 51/2 [20 + 50 \times 4]$ $= 51/2 \times 220$ $= 51 \times 110$ $= 5610$	 1 1 1 1
32. (b)	nth term of an AP $a_n = a + (n - 1)d$ Let a be the first term and d the common difference. ..... According to the question, $a_3 = 16$ and $a_7 - a_5 = 12$ $a + (3 - 1)d = 16$ $a + 2d = 16$ ..... (1) ..... Using $a_7 - a_5 = 12$ $[a + (7 - 1) d] - [a + (5 - 1) d] = 12$ $[a + 6d] - [a + 4d] = 12$ $2d = 12$ $d = 6$ ..... By substituting this in equation (1), we obtain $a + 2 \times 6 = 16$	1/2 1 $1\frac{1}{2}$

	a + 12 = 16 a = 4 ..... Therefore, A.P. will be 4, 4 + 6, 4 + 2 × 6, 4 + 3 × 6, ... Hence, the sequence will be 4, 10, 16, 22, ...	1 1
33. (a)	**<u>Statement:Basic Proportionality Theorem</u>** Prove that if a line is drawn parallel to one side of a triangle ,the other two sides are divided in the same ratio. **…..……..** **Given:** In ΔABC, DE||BC . . . . . .  . . . . To prove: $\frac{AD}{DB} = \frac{AE}{EC}$	1 1/2 1/2 1/2

	---------- **Construction :** Draw EM ⊥ AB and DN ⊥ AC. Join B to E and C to D ---------- **Proof**: In ΔADE and ΔBDE $\frac{\textbf{Area of}\Delta\textbf{ADE}}{\textbf{Area of }\Delta\textbf{BDE}} = \frac{\frac{1}{2}\times\textbf{AD}\times\textbf{EM}}{\frac{1}{2}\times\textbf{DB}\times\textbf{EM}} = \frac{\textbf{AD}}{\textbf{DB}}$ ----------(i) ---------- In ΔADE and ΔCDE $\frac{\textbf{Area of}\Delta\textbf{ADE}}{\textbf{Area of }\Delta\textbf{CDE}} = \frac{\frac{1}{2}\times\textbf{AE}\times\textbf{DN}}{\frac{1}{2}\times\textbf{EC}\times\textbf{DN}} = \frac{\textbf{AE}}{\textbf{EC}}$ ----------(ii) ---------- Since, DE||BC [Given] ∴ ar(ΔBDE) = ar(ΔCDE) ---------- (iii) [Δs on the same base and between the same parallel sides are equal in area]	1/2 1/2 1/2 1/2

	---------- From eq. (i), (ii) and (iii) : $\frac{\mathbf{AD}}{\mathbf{DB}} = \frac{\mathbf{AE}}{\mathbf{EC}}$ Hence proved.	1/2
33. (b)	 ..... Given, ΔABC ~ ΔPQR ⇒ ∠ABC = ∠PQR (corresponding angles) --------- (1) ⇒ AB/PQ = BC/QR (corresponding sides) ..... ⇒ AB/PQ = (BC/2) / (QR/2) ⇒ AB/PQ = BD/QM (D and M are mid-points of BC and QR) ---------- (2) ..... In ΔABD and ΔPQM, ∠ABD = ∠PQM (from 1) AB/PQ = BD/QM (from 2) ⇒ ΔABD ~ ΔPQM (SAS criterion) ..... . ⇒ AB/PQ = BD/QM = AD/PM (corresponding sides) ⇒ AB/PQ = AD/PM Hence proved.	1/2 1 1 $1\frac{1}{2}$ 1

34. (a)	 Depth of each conical depression, h_1 = 1.4 cm Radius of each conical depression, r = 0.5 cm Dimensions of the cuboid are 15 cm × 10 cm × 3.5 cm …… Volume of wood in the entire pen stand = volume of the wooden cuboid - 4 × volume of the conical depression …… $= l \times b \times h - 4 \times 1/3\ \pi r^2 h_1$ …… = (15 cm × 10 cm × 3.5 cm) - (4 × 1/3 × 22/7 × 0.5 cm × 0.5 cm × 1.4 cm) …… = 525 cm^3 - 1.47 cm^3 = 523.53 cm^3 The volume of wood in the entire stand is 523.53 cm^3.	1 1 1 1 1
34. (b)	The total surface area of the cube	

	$=6\times(\text{edge})^2=6\times5\times5\ \text{cm}^2=150\ \text{cm}^2$.	1
	……	
	The surface area of the block = Total Surface Area of cube - base area of hemisphere + Curved Surface Area of hemisphere	1
	……	
	$=150-\pi r^2+2\pi r^2$	1
	$=(150+\pi r^2)\ \text{cm}^2$,	
	……	
	$=150\ \text{cm}^2+(22/7\times4.2/2\times4.2/2)\ \text{cm}^2$	1
	$=(150+13.86)\ \text{cm}^2$	
	……	
	$=163.86\ \text{cm}^2$	1

35. (a)

class interval	class-mark (x_i)	Number of children(f_i)	f_ix_i
11-13	12	7	84
13-15	14	6	84
15-17	16	9	144
17-19	18	13	234
19-21	20	f	20f
21-23	22	5	110
23-25	24	4	96
		$\sum f_i=44+f$	$\sum f_ix_i=752+20f$

1+1

……

	Mean $= \bar{X} = \frac{\sum f_i x_i}{\sum f_i}$	1/2
	.....	
	$\Rightarrow 18 = \frac{752+20f}{44+f}$	1/2
	.....	
	$\Rightarrow$ 18(44+f) =752+20f	1/2
	$\Rightarrow$ 792 +18f = 752+ 20f	
	.....	
	$\Rightarrow$792-752 = 20f -18 f	1/2
	.....	
	$\Rightarrow$40 = 2f	
	$\Rightarrow$ f = 20	1
	Hence, missing frequency f =20	
35. (b)		

Number of Cars	Frequency
0-10	7
10-20	14
20-30	13
30-40	12
40-50	20
50-60	11
60-70	15
70-80	8

From the table, it can be observed that the maximum class frequency is 20, belonging to class interval 40 − 50 — 1

Therefore, modal class = 40 − 50

.....

Class size, h = 10

Lower limit of modal class, l = 40

Frequency of modal class, f_1 = 20 — 1

Frequency of class preceding modal class, f_o = 12

	Frequency of class succeeding the modal class, $f_2 = 11$ …… Mode = $l + [(f_1 - f_0)/(2f_1 - f_0 - f_2)] \times h$	1
	…… $= 40 + [(20 - 12)/(2 \times 20 - 12 - 11)] \times 10$	1/2
	…… $= 40 + [8/(40 - 23)] \times 10$ $= 40 + (8/17) \times 10$ $= 40 + 4.705$	1
	…… $= 44.705$ ≈ 44.7 Hence, the mode is 44.7	1/2
	Section - E	
36.	(i) Time = $\frac{\text{Distance}}{\text{Speed}}$	
	(ii)Let the usual speed of plane be x km/h New increased speed of plane= (x + 250)km/h Total distance=1500 km According to question $\frac{1500}{x} - \frac{1500}{x+250} = \frac{1}{2}$	1/2
	…… $\frac{1500(x + 250) - 1500x}{x(x+250)} = \frac{1}{2}$ $\frac{1500x + 375000 - 1500x}{x(x+250)} = \frac{1}{2}$ $X^2 + 250x = 750000$ $X^2 + 250x - 750000 = 0$	1/2


	(iii)(a) X^2 +250 x - 750000=0 X^2 +1000x - 750x- 750000=0 X(x+1000)-750(x +1000)=0 (x+1000)(x-750)=0 …… X=-1000 or x = 750 Reject x=-1000, because speed cannot be negative. Hence,usual speed of plane is 750km/h.	1 1
	(iii)(b) X^2 +250 x - 750000=0 X^2 +1000x - 750x- 750000=0 X(x+1000)-750(x +1000)=0 (x+1000)(x-750)=0 …… X=-1000 or x = 750 Reject x=-1000, because speed cannot be negative. Hence,new speed of plane is x+250= 750+250=1000km/h.	1 1
37.	(i) Since,radius at a point of contact is perpendicular to tangent. ∴ By Pythagoras theorem, we have PA = $\sqrt{PS^2 + AS^2}$=$\sqrt{12^2 + 5^2}$ =$\sqrt{169}$ = 13 cm …… (ii)one common tangent can be drawn when two circles touch externally. …… (iii)(a) By Pythagoras theorem, we have BQ = $\sqrt{TQ^2 + TQ^2}$=$\sqrt{3^2 + 4^2}$ =$\sqrt{25}$ = 5 cm …… QY= BQ- BY = 5-4 =1 cm ……	 1 1 1 1

	(iii) (b) PK= PA + AK = 13 + 5 = 18 cm …… XY = XK + KY= 10 +8 = 18 cm	1 1
38.	(i) Total no. of fish in the aquarium = 13+18+12+11= 54 Number of male fish in the aquarium = 36 ∴Number of female fish in the aquarium =54- 36 =18 So,probability of selecting a female fish= $\frac{\text{no. of favourable outcomes}}{\text{total no. of possible outcomes}} = \frac{18}{54} = \frac{1}{3}$ …… (ii)The probability of selecting a flowerhorn fish= $\frac{\text{no. of favourable outcomes}}{\text{total no. of possible outcomes}} = \frac{18}{54} = \frac{1}{3}$ …… (iii) (a) The probability of selecting a koi fish =$\frac{\text{no. of favourable outcomes}}{\text{total no. of possible outcomes}} = \frac{12}{54} = \frac{2}{9}$ …… P(selecting a guppy fish) =$\frac{\text{no. of favourable outcomes}}{\text{total no. of possible outcomes}} = \frac{13}{54}$	1 1 1 1

	(iii) (b)Total no. of angel fish and flowerhorn fish= 18 +11 =29 P (selecting either angel fish or flowerhorn fish)= $\frac{29}{54}$ …… P (selecting neither angel fish nor flowerhorn fish) = =1- P (selecting either angel fish or flowerhorn fish) = 1- $\frac{29}{54} = \frac{25}{54}$	1 1

MARKING SCHEME BSEH PRACTICE PAPER 2, 10TH गणित (आधार) , March2025 (हिंदी माध्यम)		
Q. no.	**Expected solutions**	**marks**
	खण्ड-क	
1	(d)60	1
2	(d) 3 से अधिक	1
3	(c)$(x+2)(x-1)=x^2-2x-3$	1
4	(c)3 इकाई	1
5	(a) -12	1
6	(a) 50°	1
7	(d) 55°	1
8	(b) $\frac{b}{\sqrt{a^2+b^2}}$	
9	(a)60^0	1
10	(b) $10\sqrt{2}$	1
11	(d) 3	1
12	(a) $\frac{1}{5}$	1
13	अपरिमेय संख्या	1
14	$\sqrt{119}$ cm	1
15	tanθ =a b	1
16	$\frac{1}{2}$	1
17	$\frac{77}{2}$ cm^2 or $\frac{49\pi}{4}$ cm^2	1
18	असत्य	1
19	(a) अभिकथन (A) और तर्क (R) दोनों सही हैं और तर्क (R), अभिकथन (A) की सही व्याख्या करता है।	1

20	(b) अभिकथन (A) और तर्क (R) दोनों सही हैं और तर्क (R), अभिकथन(A) की सही व्याख्या नहीं है।	1
	खण्ड -ख	
21. (a)	x/2 + 2y/3 = -1 3x + 4y = -6 ..... (i) ..... x-y/3 = 3 3x – y = 9 ..... (ii) ..... जब समीकरण (ii) को समीकरण (i) से घटाया जाता है तो हमें प्राप्त होता है, 5y = -15 y = -3 .....(iii) ..... जब समीकरण (iii) को (i) में प्रतिस्थापित किया जाता है तो हमें प्राप्त होता है, 3x – 12 = -6 3x = 6 x = 2 अत:, x = 2 , y = -3	1/2 1/2 1/2 1/2
21. (b)	एक आयत के गुण का उपयोग करते हुए, हम जानते हैं कि, लंबाई समान हैं, i.e., CD = AB Hence, x + 3y = 13 ...(i) ..... चौड़ाई बराबर हैं, i.e., AD = BC अत:, 3x + y = 7 ...(ii) ..... समीकरण (ii) को 3 से गुणा करके और फिर समीकरण (i) से घटाने पर हमें प्राप्त होता है, 8x = 8 So, x = 1	1/2 1/2 1/2

	समीकरण(i) में x = 1 रखने पर , हमें मिलता है, y = 4 इसलिए, x और y के वांछित मान क्रमशः 1 और 4 हैं	1/2
22.	मान लीजिए (- 4, 6) AB को आंतरिक रूप से k: 1 के अनुपात में विभाजित करता है। विभाजन सूत्र का उपयोग करते हुए, हमें मिलता है $(-4, 6) = (\frac{3k-6}{k+1}, \frac{-8k+10}{k+1})$ ..... So, $-4 = \frac{3k-6}{k+1}$ ..... $\Rightarrow -4k - 4 = 3k - 6$ $\Rightarrow 7k = 2$ $\Rightarrow, k : 1 = 2 : 7$ हम y-निर्देशांक की भी जांच कर सकते हैं। अतः, बिंदु (- 4, 6), बिंदु A(- 6, 10)और बिंदु B(3, – 8) को मिलाने वाले रेखाखंड को 2:7 के अनुपात में विभाजित करता है ।	1 1/2 1/2
23.	 ΔPBC और ΔPDE में, ∠BPC = ∠EPD [शिर्षाभिमुख कोण] PB/PD = 5/10 = ½ ... (i) PC/PE = 6/12 = ½ ... (ii) ..... समीकरण (i) और (ii) से,	1

	हमें मिलता है, PB/PD = PC/PE क्योंकि,ΔPBC का ∠BPC= ΔPDE का ∠EPD तथा उनकी सम्मिलित भुजाएं भी समानुपाती हैं …… ∴ SAS समरूपता कसौटी द्वारा ΔPBC ~ ΔPDE	1/2 1/2
24. (b)	हम जानते है कि cos 60° = 1/2 sec 30° = 2/√3 tan 45° = 1 sin 30° = 1/2 cos 30° = √3/2 …… उपरोक्त मानों को दिए गए प्रश्न में रखने पर $(5\cos^2 60° + 4\sec^2 30° – \tan^2 45°)/(\sin^2 30° + \cos^2 30°)$ $= \{5(1/2)^2+4(2/\sqrt{3})^2-1\}/(1/2)^2+(\sqrt{3}/2)^2$ …… = (5/4+16/3-1)/(1/4+3/4) ={ (15+64-12)/12}/(4/4) …… = 67/12	1/2 1/2 1/2 1/2
24. (a)	LHS = $\sqrt{\frac{1+\sin A}{1-\sin A}} =$	

	$= \sqrt{\frac{1+sinA}{1-sinA} \times \frac{1+sinA}{1+sinA}}$ ……	1/2
	$=\frac{1+\text{sinA}}{\sqrt{1-\sin^2 \text{A}}}$ ……	1/2
	$= \frac{1+\text{sinA}}{\sqrt{\cos^2 \text{A}}}$ ……	1/2
	$= \frac{1+\text{sinA}}{\text{cosA}}$ = secA +tanA = RHS	1/2
25.	मिनट की सुई द्वारा 60 मिनट में तय किया गया क्षेत्रफल =मिनट की सूई की लंबाई के बराबर त्रिज्या वाले वृत्त का क्षेत्रफल=πr^2 ……	1/2
	मिनट की सुई द्वारा 1 मिनट में तय किया गया क्षेत्रफल = $\pi r^2/60$ ……	1/2
	अत:,मिनट की सुई द्वारा 5 मिनट में तय किया गया क्षेत्रफल = $(\pi r^2/60) \times 5 = \pi r^2/12$ [∵मिनट की सुईकी लम्बाई (r) = 14 cm] ……	1/2


	= 1/12 × 22/7 × 14 × 14 cm^2 = 154/3 cm^2	1/2
	खण्ड -ग	
26.	मान लीजिए,यदि संभव हो तो ,$\sqrt{2}$ एक परिमेय संख्या है । ---------- ---------- $\therefore \sqrt{2} = \frac{p}{q}$, जहाँ p और q सह अभाज्य पूर्णांक हैं तथा $q \neq 0$. ---------- $\Rightarrow 2 = \frac{p^2}{q^2}$ $\Rightarrow p^2 = 2q^2$ .....(i) $\Rightarrow$ 2 , p^2को विभाजित करता है $\Rightarrow$ 2 , p को भी विभाजित करता है । ---------- माना p = 2m,.....(ii) जहाँ m कोई पूर्णांक है । $\Rightarrow p^2 = 4m^2$.....(iii) ---------- (i) और(iii) से $2q^2 = 4m2$ $\Rightarrow q^2 = 2m^2$ $\Rightarrow$ 2 , q^2को विभाजित करता है $\Rightarrow$ 2 , q को भी विभाजित करता है । $\Rightarrow$ q = 2n.....(iv) ---------- (ii) और (iv) से, p और q का उभयनिष्ठ गुणनखंड 2 है। $\therefore$ p और q सह-अभाज्य पूर्णांक नहीं हैं अतः हमारी कल्पना गलत है। $\therefore \sqrt{2}$ एक अपरिमेय संख्या है।	1/2 1/2 1/2 1/2 1/2 1/2

27.	$6x^2-3-7x=6x^2-7x-3=0$ $\Rightarrow 6x^2+2x-9x-3=0$ $\Rightarrow 2x(3x+1)-3(3x+1)=0$ $\Rightarrow (2x-3)(3x+1)=0$ शून्यकα,β= 3/2,−1/3 ……	1
	$\alpha+\beta=-b/a \Rightarrow (3/2)+(-1/3)=7/6=-(-7)6=-b/a$ ……	1
	$\alpha\beta=c/a \Rightarrow (3/2)(-1/3)=-1/2=-3/6=c/a$ अत: सिद्ध हुआ।	1
28. (a)	माना कि राहुल की आयु x वर्ष है और उसके पुत्र की आयु y वर्ष है। ……	1/2
	पाँच वर्ष बाद (बाद में), x+ 5 = 3 (y + 5) $\Rightarrow$x+ 5 = 3 y + 15 $\Rightarrow$x - 3 y = 10.....(1) ……	1/2
	साथ ही, पांच साल पूर्व(पहले), x-5 = 7 (y – 5) $\Rightarrow$ x-5 = 7y – 35 $\Rightarrow$x-7y = -30.....(2) ……	1/2
	समीकरण (2) को (1) से घटाने पर, x - 3 y -x +7y = 10+ 30	

	(∵समीकरण(2) का चिन्ह बदलता है) 4y = 40 ⇒y = 10 …… समीकरण(1) में $y = 10$ रखने पर। x - 3(10) = 10 ⇒x - 30 = 10 ⇒x = 40 …… अत: राहुल की वर्तमान आयु=x=40 वर्ष और राहुल के पुत्र की वर्तमान आयु=y=10 वर्ष.	1/2 1/2 1/2
28. (b)	माना बड़ा कोण = x छोटा कोण = y चूँकि दोनों कोण संपूरक हैं, ∴ x + y = 180 ⇒ x = 180 - y .... (i) …… अंतर 18 डिग्री है. ∴ x - y = 18 ⇒ x = 18 + y .... (i) …… समीकरण (i) में x का मान रखने पर हमें मिलता है, ⇒ 18 + y = 180 - y ⇒ - y - y = 18 - 180 ⇒ - 2y = -162 ⇒ y= −162/−2 ⇒ y = 81 ……	 1 1/2 1/2


	समीकरण (i) में y का मान रखने पर, हमें मिलता है, x = 180 - 81 = 99 ..... इसलिए, कोण 99° और 81° हैं।	1/2 1/2
29.	हम जानते हैं कि दो बिंदुओं के बीच की दूरी, दूरी सूत्र $= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ द्वारा दी जाती है दूरी सूत्र में बिंदु P (2, - 3) और Q (10, y) के मानों को प्रतिस्थापित करने पर, हमें मिलता है ..... PQ = $\sqrt{(2 - 10)^2 + (-3 - y)^2} = 10$ PQ = $\sqrt{(-8)^2 + (3 + y)^2} = 10$ ..... दोनों तरफ वर्ग करने पर, हमें मिलता है $64 + (y + 3)^2 = 100$ ..... $(y + 3)^2 = 36$ $y + 3 = \sqrt{36}$ $y + 3 = \pm 6$ ..... y + 3 = 6 या y + 3 = - 6 इसलिए, y = 3 या - 9, y के लिए संभावित मान हैं।	1/2 1/2 1/2 1 1
30. (a)	दिया है: $\cos A + \cos^2 A = 1$ $\Rightarrow \cos A = 1 - \cos^2 A$ $\Rightarrow \cos A = \sin^2 A$ [∵ $\sin^2 A = 1 - \cos^2 A$] .....(i)	1

	…… LHS= $(\sin^2 A + \sin^4 A) = (\sin^2 A + (\sin^2 A)^2)$ …… $= (\sin^2 A + (\cos A)^2)$ [(i)का उपयोग करने पर] …… $= \sin^2 A + \cos^2 A$ $= 1 = RHS$	1/2 1 1/2
30. (b)	LHS=$(\sin A+\operatorname{cosec} A)^2+(\cos A+\sec A)^2$ $=\sin^2 A+\operatorname{cosec}^2 A+2\sin A\operatorname{cosec} A+\cos^2 A+\sec^2 A+2\cos A\sec A$ …… $=\sin^2 A+\cos^2 A+\operatorname{cosec}^2 A+\sec^2 A+2\sin A\times 1/\sin A+2\cos A\times 1/\cos A$ [∵cosecA=1/sinA and secA=1/cosA] …… $=1+\operatorname{cosec}^2 A+\sec^2 A+2+2$ [∵$\sin^2 A+\cos^2 A=1$] …… $=5+(1+\cot^2 A)+(1+\tan^2 A)$ [∵$1+\tan^2 A=\sec^2 A$ and $1+\cot^2 A=\operatorname{cosec}^2 A$] $=7+\tan^2 A+\cot^2 A=$ RHS	1/2 1 1/2 1

31.	 …… प्रेक्षक की ऊंचाई DE = 1.5 मीटर दी गई है AB = DE=1.5 m माना BC = h चिमनी की ऊँचाई है अत: AC = (h – 1.5) m प्रेक्षक और चिमनी के बीच की दूरी AD = BE = 28.5 मीटर है …… समकोण △CAD में ,θ=45^{o} tan 45^{o}=AC/AD ⇒1=(h−1.5)/28.5 …… ⇒28.5=h−1.5 ⇒h=28.5+1.5= 30 m अतः चिमनी की ऊंचाई 30 मीटर है।	1/2 1/2 1 1

खण्ड-घ

32. (a)	दिया है: दूसरा पद $a_2 = 14$ तीसरा पद $a_3 = 18$ सार्व अंतर $d = a_3 - a_2 = 18 - 14 = 4$ ..... हम जानते है कि AP का nवाँ पद होता है: $a_n = a + (n - 1)d$ $a_2 = a + d$ $14 = a + 4$ $a = 10$ ..... AP के n पदों का योगफल होता है :$S_n = n/2 [2a + (n - 1) d]$ ..... $\therefore S_{51} = 51/2 [2 \times 10 + (51 - 1) 4]$ ..... $= 51/2 [20 + 50 \times 4]$ $= 51/2 \times 220$ $= 51 \times 110$ $= 5610$	1 1 1 1 1
32. (b)	AP का nवाँ पद होता है: $a_n = a + (n - 1)d$ जहाँ a प्रथम पद तथा d सार्व अंतर होता है ..... प्रश्न के अनुसार , $a_3 = 16$ और $a_7 - a_5 = 12$ $a + (3 - 1)d = 16$ $a + 2d = 16$ ..... (1) ..... $a_7 - a_5 = 12$ का उपयोग करते हुए $[a + (7 - 1) d] - [a + (5 - 1) d] = 12$	1/2 1

	[a + 6d] - [a + 4d] = 12 2d = 12 d = 6 ..... इसे समीकरण (1) में रखने पर हमें प्राप्त होता है : a + 2 × 6 = 16 a + 12 = 16 a = 4 ..... अत:अभिष्ठ AP होगी : 4, 4 + 6, 4 + 2 × 6, 4 + 3 × 6, ... या 4, 10, 16, 22, ...	$1\frac{1}{2}$ 1 1
33. (a)	**<u>कथन: आधारभूत समानुपातिकता प्रमेय</u>** **<u>सिद्ध कीजिए कि यदि किसी त्रिभुज की एक भुजा के समानांतर एक रेखा खींची जाए, तो अन्य दो भुजाएँ समान अनुपात में विभाजित हो जाती हैं।</u>** **…..** **दिया है:** ΔABC में, DE||BC **.....**  	 1/2 1/2

.....

सिद्ध करना है : $\frac{AD}{DB} = \frac{AE}{EC}$ 1/2

रचना : EM⊥AB तथा DN⊥AC खींचिए। B को E से तथा C को D से मिलाइये । 1/2

प्रमाण: ΔADE तथा ΔBDE में

$$\frac{\text{ΔADE का क्षेत्रफल}}{\text{ΔBDE का क्षेत्रफल}} = \frac{\frac{1}{2}\times AD\times EM}{\frac{1}{2}\times DB\times EM} = \frac{AD}{DB} \text{----------(i)}$$ 1/2



ΔADE तथा ΔCDE में


$$\frac{\text{ΔADE का क्षेत्रफल}}{\text{ΔCDE का क्षेत्रफल}} = \frac{\frac{1}{2}\times AE\times DN}{\frac{1}{2}\times EC\times DN} = \frac{AE}{EC} \text{----------(ii)}$$ 1/2

क्योंकि DE||BC [दिया है]

∴ (ΔBDE) का क्षेत्रफल = (ΔCDE)का क्षेत्रफल ---------- (iii) 1/2

[∵एक ही आधार पर और एक ही समानांतर भुजाओं के बीच बनी त्रिभुजों का क्षेत्रफल बराबर होता है]

	---------- समीकरण (i), (ii) और (iii)से : $\frac{AD}{DB} = \frac{AE}{EC}$ यही सिद्ध करना था।	1/2
33. (b)	  ..... दिया है : ΔABC ~ ΔPQR ⇒ ∠ABC = ∠PQR (संगत कोण) --------- (1) ⇒ AB/PQ = BC/QR (संगत भुजाएं) ..... ⇒ AB/PQ = (BC/2) / (QR/2) ⇒ AB/PQ = BD/QM (∵D और M ,BC तथाQR के मध्य बिंदु हैं।) ---------- (2) ..... ΔABD तथा ΔPQM में, ∠ABD = ∠PQM ((1)से) AB/PQ = BD/QM ((2)से) ⇒ ΔABD ~ ΔPQM (SAS कसौटी द्वारा) ..... ⇒ AB/PQ = BD/QM = AD/PM (संगत भुजाएं) ⇒ AB/PQ = AD/PM यही सिद्ध करना था।	1/2 1 1 $1\frac{1}{2}$ 1

34. (a)	 प्रत्येक शंक्वाकार गड्ढे की गहराई, h_1 =1.4 सेमी प्रत्येक शंक्वाकार गड्ढे की त्रिज्या, r = 0.5 सेमी घनाभ का आयाम 15 सेमी × 10 सेमी × 3.5 सेमी है ..... पूरे पेन स्टैंड में लकड़ी का आयतन = लकड़ी के घनाभ का आयतन - 4 × शंक्वाकार गड्ढे का आयतन ..... = l × b × h - 4 × 1/3 $\pi r^2 h_1$ ..... = (15 cm × 10 cm × 3.5 cm) - (4 × 1/3 × 22/7 × 0.5 cm × 0.5 cm × 1.4 cm) ..... = 525 cm^3 - 1.47 cm^3 = 523.53 cm^3 पूरे स्टैंड में लकड़ी का आयतन 523.53 cm^3 है।	1 1 1 1 1
34. (b)	घन का कुल पृष्ठीय क्षेत्रफल	

	$=6\times(\text{भुजा})^2=6\times5\times5\ cm^2=150\ cm^2$.	1
	……	
	ब्लॉक का पृष्ठीय क्षेत्रफल = घन का कुल पृष्ठीय क्षेत्रफल - गोलार्ध का आधार क्षेत्रफल + गोलार्ध का वक्र पृष्ठीय क्षेत्रफल	1
	……	
	$=150-\pi r^2+2\pi r^2$	
	$=(150+\pi r^2)\ cm^2$,	1
	……	
	$=150\ cm^2+(22/7\times4.2/2\times4.2/2)\ cm^2$	
	$=(150+13.86)\ cm^2$	1
	……	
	$=163.86\ cm^2$	1
35. (a)		1+1

वर्ग -अंतराल	वर्ग -चिन्ह(x_i)	बच्चों की संख्या(f_i)	f_ix_i
11-13	12	7	84
13-15	14	6	84
15-17	16	9	144
17-19	18	13	234
19-21	20	f	20f
21-23	22	5	110
23-25	24	4	96
		$\sum f_i=44+f$	$\sum f_ix_i=752+20f$

……

	माध्य $= \bar{X} = \frac{\sum f_i x_i}{\sum f_i}$	1/2
	.....	
	$\Rightarrow 18 = \frac{752+20f}{44+f}$	1/2
	.....	
	$\Rightarrow$ 18(44+f) =752+20f	1/2
	$\Rightarrow$ 792 +18f = 752+ 20f	
	.....	
	$\Rightarrow$792-752 = 20f -18 f	1/2
	.....	
	$\Rightarrow$40 = 2f	
	$\Rightarrow$ f = 20	1
	अत:लुप्त बारंबारता $f = 20$	
35. (b)		

कारों की संख्या	बारंबारता
0-10	7
10-20	14
20-30	13
30-40	12
40-50	20
50-60	11
60-70	15
70-80	8

तालिका से, यह देखा जा सकता है कि अधिकतम वर्ग बारंबारता 20 है, जो वर्ग अंतराल 40 - 50 से संबंधित है।	1
इसलिए, बहुलक वर्ग = 40 – 50	
.....	
वर्ग आमाप h = 10	
बहुलक वर्ग की निचली सीमा, $l = 40$	1
बहुलक वर्ग की बारंबारता, $f_1 = 20$	

	बहुलक वर्ग से पहले वाले वर्ग कीबारंबारता, $f_0 = 12$ बहुलक वर्ग के बाद वाले वर्ग की बारंबारता, $f_2 = 11$ …… बहुलक = $l + [(f_1 - f_0)/(2f_1 - f_0 - f_2)] \times h$ …… $= 40 + [(20 - 12)/(2 \times 20 - 12 - 11)] \times 10$ …… $= 40 + [8/(40 - 23)] \times 10$ $= 40 + (8/17) \times 10$ $= 40 + 4.705$ …… $= 44.705$ ≈ 44.7 अत: बहुलक 44.7	1 1/2 1 1/2
	खण्ड-ङ	
36.	(i) समय $= \frac{\text{दूरी}}{\text{गति}}$	1
	(ii)माना विमान की सामान्य गति x किमी/घंटा है विमान की नई बढ़ी हुई गति = (x + 250) किमी/घंटा कुल दूरी=1500 किमी प्रश्न के अनुसार $\frac{1500}{x} - \frac{1500}{x+250} = \frac{1}{2}$ …… $\frac{1500(x + 250) - 1500x}{x(x+250)} = \frac{1}{2}$ $\frac{1500x + 375000 - 1500x}{x(x+250)} = \frac{1}{2}$ $X^2 + 250x = 750000$	1/2

	X^2 +250 x - 750000=0	1/2
	(iii)(a) X^2 +250 x - 750000=0 X^2 +1000x - 750x- 750000=0 X(x+1000)-750(x +1000)=0 (x+1000)(x-750)=0 …… X=-1000 अथवा x = 750 x=-1000 को अस्वीकार करें, क्योंकि गति ऋणात्मक नहीं हो सकती। अतः, विमान की सामान्य गति 750 किमी/घंटा है।	1 1
	(iii)(b) X^2 +250 x - 750000=0 X^2 +1000x - 750x- 750000=0 X(x+1000)-750(x +1000)=0 (x+1000)(x-750)=0 …… X=-1000 अथवा x = 750 x=-1000 को अस्वीकार करें, क्योंकि गति ऋणात्मक नहीं हो सकती। अतः, विमान की नई गति x+250= 750+250=1000 किमी/घंटा है।	1 1
37.	(i) चूँकि, संपर्क बिंदु पर त्रिज्या स्पर्श रेखा के लंबवत होती है। ∴ पाइथागोरस प्रमेय से PA = $\sqrt{PS^2 + AS^2}$=$\sqrt{12^2 + 5^2}$ =$\sqrt{169}$ = 13 cm …… (ii)जब दो वृत्त बाह्य रूप से स्पर्श करते हैं तो एक उभयनिष्ठ स्पर्शरेखा खींची जा सकती है। …… (iii)(a) पाइथागोरस प्रमेय से BQ = $\sqrt{TQ^2 + TQ^2}$=$\sqrt{3^2 + 4^2}$ =$\sqrt{25}$ = 5 cm	1 1 1

	.....	
	QY= BQ- BY = 5-4 =1 cm	1
	.....	
	(iii) (b) PK= PA + AK = 13 + 5 = 18 cm	1
	.....	
	XY = XK + KY= 10 +8 = 18 cm	1
38.	(i) एक्वेरियम में कुल मछलियाँ = 13+18+12+11= 54 एक्वेरियम में कुल नर मछलियाँ = 36 ∴एक्वेरियम में मादा मछलियाँ =54- 36 =18 ∴मादा मछली के चयन की प्रायिकता = $\frac{\text{अनुकूल परिणामोंकी संख्या}}{\text{कुल संभावित परिणामों की संख्या}} = \frac{18}{54} = \frac{1}{3}$	
	.....	
	(ii)फ्लावरहॉर्न मछली को चुनने की प्रायिकता = $\frac{\text{अनुकूल परिणामोंकी संख्या}}{\text{कुल संभावित परिणामों की संख्या}} = \frac{18}{54} = \frac{1}{3}$	1
	.....	
	(iii) (a) 'कोई' मछली के चुनने की प्रायिकता= $\frac{\text{अनुकूल परिणामोंकी संख्या}}{\text{कुल संभावित परिणामों की संख्या}} = \frac{12}{54} = \frac{2}{9}$	1

	.....	
	गप्पी मछली के चुनने की प्रायिकता= $\frac{\text{अनुकूल परिणामोंकी संख्या}}{\text{कुल संभावित परिणामों की संख्या}} = \frac{13}{54}$	1
	(iii) (b)एंजल और फ्लावरहॉर्न मछलियोंकी कुल संख्या= 18 +11 =29	
	एंजल मछली या फ्लावरहॉर्न मछली को चुनने की प्रायिकता= $\frac{29}{54}$	1
	.....	
	न ही एंजल मछली और न ही फ्लावरहॉर्न मछली को चुनने की प्रायिकता=	
	=1- एंजल मछली या फ्लावरहॉर्न मछली को चुनने की क्रमशः प्रायिकता	1
	= 1- $\frac{29}{54} = \frac{25}{54}$	

Sample Papers

- CBSE Sample Papers
- ICSE / ISC Board Sample Papers
- AP Board Sample Papers
- Assam Board Sample Papers
- Bihar Board Sample Papers
- Chhattisgarh Board Sample Papers
- Goa Board Sample Papers
- Gujarat Board Sample Papers
- Haryana Board Sample Papers
- HP Board Sample Papers
- J&K State Board Sample Papers
- Jharkhand Board Sample Papers
- Karnataka Board Sample Papers
- Kerala Board Sample Papers

Sample Papers

- MP Board Sample Papers
- Maharashtra Board Sample Papers
- Manipur Board Sample Papers
- Mizoram Board Sample Papers
- Orissa Board Sample Papers
- Punjab Board Sample Papers
- Rajasthan Board Sample Papers
- Tamil Nadu Board Sample Papers
- Telangana State Board Sample Papers
- Tripura Board Sample Papers
- UP Board Sample Papers
- Uttarakhand Board Sample Papers
- West Bengal Board Sample Papers

