NCERT SOLUTIONS

CLASS - 12th



aglasem.com

Class : 12th
Subject : Chemistry
Chapter : 10
Chapter Name : Haloalkanes and Haloarenes

Q10.1 Write structures of the following compounds:
(i) 2-Chloro-3-methylpentane
(ii) I-Chloro-4-ethylcyclohexane
(iii) 4-tert. Butyl-3-iodoheptane
(iv) 1,4-Dibromobut-2-ene
(v) I-3romo-4-sec. butyl-2-methylbenzene

Answer.

(i)



2-Chloro-3-methyl pentane

(ii)



1-Chloro-4-ethylcyclohexane

(iii)



4- tert-Butyl-3-iodoheptane

(iv)



1, 4-Dibromobut-2-ene

(v)



1-Bromo-4-sec-butyl-2-methylbenzene

Page : 293 , Block Name : Intext Questions

Q10.2 Why is sulphuric acid not used during the reaction of alcohols with KI?

Answer. In the presence of sulphuric acid H_2SO_4, KI produces HI

$$2KI + H_2SO_4 \longrightarrow 2KHSO_4 + 2HI$$

Since H_2SO_4 is an oxidizing agent, it oxidizes HI (produced in the reaction to 12).

$$2HI + H_2SO_4 \longrightarrow I_2 + SO_2 + H_2O$$

As a result, the reaction between alcohol and HI to produce alkyl iodide cannot occur. Therefore, sulphuric acid is not used during the reaction of alcohols with KI. Instead, a non-oxidizing acid such as H_3PO_4 is used.

Page : 297 , Block Name : Intext Questions

Q10.3 Write structures of different dihalogen derivatives of propane.

Answer. There are four different dihalogen derivatives of propane. The structures of these derivatives are shown below.

(i)



1, 1-Dibromopropane

(ii)



2, 2-Dibromopropane

(iii)



1, 2-Dibromopropane

(iv)



1, 3-Dibromopropane

Page : 297 , Block Name : Intext Questions

Q10.4 Among the isomeric alkanes of molecular formula C_5H_{12}, identify the one that on photochemical chlorination yields
(i) A single monochloride.
(ii) Three isomeric monochlorides.
(iii) four isomeric monochlorides.

Answer. (i) To have a single monochloride, there should be only one type of H-atom in the isomer of the alkane of the molecular formula C_5H_{12}. This is because, replacement of any H-atom leads to the formation of the same product. The isomer is neopentane.



(ii) To have three isomeric monochlorides, the isomer of the alkane of the molecular formula C_5H_{12} should contain three different types of H-atoms. Therefore, the isomer is n-pentane. It can be observed that there are three types of H atoms labelled as a, b and cin n-pentane.

$$CH_3 - CH_2 - CH_2 - CH_2 - CH_3$$

n -Pentane

(iii) To have four isomeric monochlorides, the isomer of the alkane of the molecular formula C_5H_{12} should contain four different types of H-atoms. Therefore, the isomer is 2- methylbutane. It can be observed that there are four types of H-atoms labelled as a, b, c, and d in 2-methylbutane.

$$CH_3 - \underset{\displaystyle CH_3}{\underset{|}{CH}} - CH_2 - CH_3$$

Page : 297 , Block Name : Intext Questions

Q10.5 Draw the structures of major monohalo products in each of the following reactions:

(i)



(ii)



(iii)



(iv)



(v)



(vi)



Answer.

(i)



Cyclohexanol — Chlorocyclohexane

(ii)



4 – Ethylnitrobenzene — 4 – (1 – Bromoethyl) nitrobenzene

(iii)



4 – Hydroxymethylphenol — 4 – Chloromethylphenol

(iv)



1 – Methylcyclohexene — 1 – Iodo – 1 – methylcyclohexane

(v)



Bromoethane — Iodoethane

(vi)



Cyclohexene — 3 – Bromocyclohexene

Page : 297 , Block Name : Intext Questions

Q10.6 Arrange each set of compounds in order of increasing boiling points.
(i) Bromomethane, Bromoform, Chloromethane, Dibromomethane.
(ii) I-Chloropropane, Isopropyl chloride, I-Chlorobutane.

Answer. (i)



Chloromethane — Bromomethane — Dibromomethane — Bromoform

For alkyl halides containing the same alkyl group, the boiling point increases with an increase in

the atomic mass of the halogen atom.
Since the atomic mass of Br is greater than that of Cl, the boiling point of bromomethane is higher than that of chloromethane. Further, for alkyl halides containing the same alkyl group, the boiling point increases with an increase in the number of halides. Therefore, the boiling point of Dibromomethane is higher than that of chloromethane and bromomethane, but lower than that of bromoform. Hence, the given set of compounds can be arranged in the order of their increasing boiling points as:
Chloromethane < Bromomethane < Dibromomethane < Bromoform.
(ii)



For alkyl halides containing the same halide, the boiling point increases with an increase in the size of the alkyl group. Thus, the boiling point of I-chlorobutane is higher than that of isopropyl chloride and I-chloropropane. Further, the boiling point decreases with an increase in branching in the chain. Thus, the boiling point of isopropyl alcohol is lower than that of I-chloropropane. Hence, the given set of compounds can be arranged in the increasing order of their boiling points as:
Isopropyl chloride < I-Chloropropane < I-Chlorobutane

Page : 299 , Block Name : Intext Questions

Q10.7 Which alkyl halide from the following pairs would you expect to react more rapidly by an S_N2 mechanism? Explain your answer.

(i)



(ii)



(iii)



Answer. (i)



2-bromobutane is a 2° alkylhalide whereas I-bromobutane is a 1° alkyl halide. The approaching of

nucleophile is more hindered in 2-bromobutane than in I-bromobutane. Therefore, I-bromobutane reacts more rapidly than 2-bromobutane by an S_N2 mechanism.

(ii)



2-3romobutane is 2° alkylhalide whereas 2-bromo-2-methylpropane is 30 alkyl halide. Therefore, greater numbers of substituents are present in 3° alkyl halide than in 2° alkyl halide to hinder the approaching nucleophile. Hence, 2-bromobutane reacts more rapidly than 2-bromo-2-methylpropane by an S_N2 mechanism.

(iii)



Both the alkyl halides are primary. However, the substituent —CH, is at a greater distance to the carbon atom linked to Br in I-bromo-3-methylbutane than in I-bromo-2-methylbutane. Therefore, the approaching nucleophile is less hindered in case of the former than in case of the latter. Hence, the former reacts faster than the latter by $S_N2$mechanism.

Page : 316 , Block Name : Intext Questions

Q10.8 In the following pairs of halogen compounds, which compound undergoes faster S_N1 reaction?



Answer. (i)



S_N1 reaction proceeds via the formation of carbocation. The alkyl halide (I) is 3° while (11) is 2°. Therefore, (I) forms 3° carbocation while (II) forms 2° carbocation. Greater the stability of the carbocation, faster is the rate of S_N1 reaction. Since 30 carbocation is more stable than 20 carbocation. (I), i.e. 2—chloro-2-methylpropane, undergoes faster S_N1 reaction than (II) i.e., 3-

chloropentane.
(ii)



The alkyl halide (I) is 2° while (II) is 1°. 2° carbocation is more stable than 1° carbocation. Therefore, (I), 2—chloroheptane, undergoes faster S_{N1} reaction than (II), 1 chlorohexane.

Page : 316 , Block Name : Intext Questions

Q10.9

Identify A, B, C, D, E, R and R^1 in the following:



Answer.



Since D of D_2O gets attached to the carbon atom to which MgBr is attached, C is



Therefore, the compound R – Br is





When an alkyl halide is treated with Na in the presence of ether, a hydrocarbon containing double the number of carbon atoms as present in the original halide is
obtained as product. This is known as Wurtz reaction. Therefore, the halide, R^1—X, is



tert – Butylhalide

Therefore, compound D is



tert – Bulytmagnesiumbromide

And, compound E is



2 – Methylpropane



2, 2, 3, 3 – tetramethylbutane



(D)

(E)

Page : 316 , Block Name : Intext Questions

Q10.1 Name the following halides according to IUPAC system and classify them as alkyl, allyl, benzyl (primary, secondary, tertian'), vinyl or aryl halides:

(i)

$(CH_3)_2CHCH(Cl)CH_3$

(ii)

$CH_3CH_2CH\,(CH_3)\,CH\,(C_2H_5)\,Cl$

(iii)

$CH_3CH_2C(CH_3)_2CH_2I$

(iv)

$(CH_3)_3CCH_2CH(Br)C_6H_5$

(v)
$CH_3CH\left(CH_3\right)CH(Br)CH_3$
(vi)
$CH_3C(C_2H_5)_2CH_2Br$
(vii)
$CH_3C(Cl)\left(C_2H_s\right)CH_2CH_3$
(viii)
$CH_3CH=C(Cl)CH_2CH(CH_3)_2$
(ix)
$CH_3CH=CHC(Br)(CH_3)_2$
(x)
$p-ClC_6H_4CH_2CH(CH_3)_2$
(xi)
$m-ClCH_2C_6H_4CH_2C(CH_3)_3$
(xii)
$o-Br-C_6H_4CH\left(CH_3\right)CH_2CH_3$

Answer.

(i)



2–Chloro–3–methylbutane

(Secondary alkyl halide)

(ii)



3–Chloro–4–methyhexane

(Secondary alkyl halide)

(iii)



1–Iodo–2, 2 –dimethylbutane

(Primary alkyl halide)

(iv)



1–Bromo–3, 3–dimethyl–1–phenylbutane

(Secondary benzyl halide)

(v)



2–Bromo–3–methylbutane

(Secondary alkyl halide)

(vi)



1–Bromo–2–ethyl–2–methylbutane

(Primary alkyl halide)

(vii)



3–Chloro–3–methylpentane

(Tertiary alkyl halide)

(viii)



3–Chloro–5–methylhex–2–ene

(Vinyl halide)

(ix)



4–Bromo–4–methylpent–2–ene

(Allyl halide)

(x)



1−Chloro−4−(2−methylpropyl) benzene

(Aryl halide)

(xi)



1−Chloromethyl−3−(2, 2−dimethylpropyl) benzene

(Primary benzyl halide)

(xii)



1−Bromo−2−(1−methylpropyl) benzene

(Aryl halide)

Page : 319 , Block Name : Exercise

Q10.2 Give the IUPAC names of the following compounds:

(i)

$\mathrm{CH_3CH(Cl)CH(Br)CH_3}$

(ii)

$\mathrm{CHF_2CBrClF}$

(iii)

$\mathrm{ClCH_2C \equiv CCH_2Br}$

(iv)

$\mathrm{(CCl_3)_3CCl}$

(v)

$\mathrm{CH_3C(\mathit{p}-ClC_6H_4)_2CH(Br)CH_3}$

(vi)

$\mathrm{(CH_3)_3CCH = CClC_6H_4I-\mathit{p}}$

Answer.

(i)



2–Bromo–3–chlorobutane

(ii)



1–Bromo–1–chloro–1, 2, 2–trifluoroethane

(iii)



1–Bromo–4–chlorobut–2–yne

(iv)



2–(Trichloromethyl)–1,1,1,2,3,3,3–heptachloropropane

(v)



2–Bromo–3, 3–bis(4–chlorophenyl) butane

(vi)



1–chloro–1–(4–iodophenyl)–3, 3–dimethylbut–1–ene

Page : 319 , Block Name : Exercise

Q10.3 Write the structures of the following organic halogen compounds.
(i)
$2-$ Chloro- 3-methylpentane
(ii)
p-Bromochlorobenzene
(iii)
$1-$ Chloro $-$ 4-ethylcyclohexane
(iv)
$2-(2-$ Chlorophenyl $)-1$-iodooctane
(v)
perfluorobenzene
(vi)
4-tert-Butyl-3-iodoheptane
(vii)
1-Bromo-4-sec-butyl-2-methylbenzene
(viii)
1,4-Dibromobut-2-ene

Answer.

(i)



2-Chloro-3-methylpentane

(ii)



p-Bromochlorobenzene

(iii)



1-Chloro-4-ethylcyclohexane

(iv)



2-(2-Chlorophenyl)-1-iodooctane

(v)



Perfluorobenzene

(vi)



4-Tert-Butyl-3-iodoheptane

(vii)



1-Bromo-4-sec-butyl-2-methylbenzene

(viii)



1,4-Dibromobut-2-ene

Page : 319 , Block Name : Exercise

Q10.4 Which one of the following has the highest dipole moment?

(i) CH_2Cl_2

(ii) $CHCl_3$

(iii) CCl_4

Answer.

(i)



Dichlormethane (CH_2Cl_2)

μ = 1.60D

(ii)



Chloroform ($CHCl_3$)

μ = 1.08D

(iii)



Carbon tetrachloride (CCl_4)

μ = 0D

CCl_4 is a symmetrical molecule. Therefore, the dipole moments cancel each other. Hence, its resultant dipole moment is zero.
As shown in the above figure, in $CHCl_3$, the resultant of dipole moments of two C Cl bonds is opposed by the resultant of dipole moments of one C—H bond and one C—CI bond. Since the resultant of one bond and one C—CI bond dipole moments is smaller than two C—CI bonds, the opposition is to a small extent. As a result, $CHCl_3$ has a small dipole moment of 1.08 D.
On the other hand, in case of CH_2Cl_2, the resultant of the dipole moments of two C—CI bonds is strengthened by the resultant of the dipole moments of two C—H bonds. As a result, CH_2Cl_2 has a higher dipole moment of 1.60 D than $CHCl_3$ i.e., CH_2Cl_2 has the highest dipole moment.
Hence, the given compounds can be arranged in the increasing order of their dipole moments as:
$CCl_4 < CHCl_3 < CH_2Cl_2$

Page : 319 , Block Name : Exercise

Q10.5 A hydrocarbon C_5H_{10} does not react with chlorine in dark but gives a single monochloro compound C_5H_9Cl in bright sunlight. Identify the hydrocarbon.

Answer. A hydrocarbon with the molecular formula, C_5H_{10} belongs to the group with a general molecular formula C_nH_{2n}. Therefore, it may either be an alkene or a cycloalkane.
Since hydrocarbon does not react with chlorine in the dark, it cannot be an alkene. Thus, it should be a cycloalkane. Further, the hydrocarbon gives a single monochloro compound, C_5H_9Cl by reacting with chlorine in bright sunlight. Since a single monochloro compound is formed, the hydrocarbon must contain H—atoms that are all equivalent. Also, as all H—atoms of a cycloalkane

are equivalent, the hydrocarbon must be a cycloalkane. Hence, the said compound is cyclopentane.



Cyclopentane (C_5H_{10})

The reactions involved in the question are:



Page : 319 , Block Name : Exercise

Q10.6 Write the isomers of the compound having formula C_4H_9Br.

Answer. There are four isomers of the compound having the formula C_4H_9Br These isomers are given below.

(a)



1–Bromobutane

(b)



2–Bromobutane

(c)



1–Bromo–2–methylpropane

(d)



2–Bromo–2–methylpropane

Page : 319 , Block Name : Exercise

Q10.7 Write the equations for the preparation of I—iodobutane from
(i) 1-butanol
(ii) 1-chlorobutane
(iii) but-1-ene.

Answer.

(i)



(ii)



(iii)



Page : 319 , Block Name : Exercise

Q10.8 What are ambident nucleophiles? Explain with an example.

Answer. Ambident nucleophiles are nucleophiles having two nucleophilic sites. Thus, ambident nucleophiles have two sites through which they can attack. For example, nitrite ion is an ambident nucleophile.



Nitrite ion can attack through oxygen resulting in the formation of alkyl nitrites. Also, it can attack through nitrogen resulting in the formation of nitroalkanes.



Page : 319 , Block Name : Exercise

Q10.9 Which compound in each of the following pairs will react faster in S_N2 reaction with OH ?
(i) CH_3Br or CH_3I
(ii) $(CH_3)_3CCl$ or CH_3Cl

Answer. (i) In the S_N2 mechanism, the reactivity of halides for the same alkyl group increases in the order. This happens because as the size increases, the halide ion becomes a better leaving group.
$R - F << R - Cl < R - Br < R - I$
Therefore, CH_3I will react faster than CH_3Br in S_N2 reactions with OH^-.
(ii)



The S_N2 mechanism involves the attack of the nucleophile at the atom bearing the leaving group. But, in case of $(CH_3)_3CCl$, the attack of the nucleophile at the carbon atom is hindered because of the presence of bulky substituents on that carbon atom bearing the leaving group. On the other hand, there are no bulky substituents on the carbon atom bearing the leaving group in $(CH_3)_3CCl$. Hence, $(CH_3)_3CCl$ reacts faster than $(CH_3)_3CCl$ in S_N2 reaction with OH

Page : 320 , Block Name : Exercise

Q10.10 Predict all the alkenes that would be formed by dehydrohalogenation of the following halides with sodium ethoxide in ethanol and identify the major alkene:
(i) I-Bromo-l-methylcyclohexane
(ii) 2-Chloro-2-methylbutane
(iii) 2,2,3-Trimethyl-3-bromopentane

Answer. (i)



1–bromo–1–methylcyclohexane

In the given compound, all ß-hydrogen atoms are equivalent. Thus, dehydrohalogenation of this compound gives only one alkene.



(ii)



In the given compound, there are two different sets of equivalent ß hydrogen atoms labelled as a and b. Thus, dehydrohalogenation of the compound yields two alkenes.



Saytzeff's rule implies that in dehydrohalogenation reactions, the alkene having a greater number of alkyl groups attached to a doubly bonded carbon atoms is preferably
produced. Therefore, alkene (I) i.e., 2-methylbut-2-ene is the major product in this reaction.

(iii)



2,2,3-Trimethyl-3-bromopentane

In the given compound, there are two different sets of equivalent ß-hydrogen atoms labelled as a and b. Thus, dehydrohalogenation of the compound yields two alkenes.



According to Saytzeff's rule, in dehydrohalogenation reactions, the alkene having a greater number of alkyl groups attached to the doubly bonded carbon atom is preferably formed. Hence, alkene (I) i.e., 3,4,4-trimethylpent-2-ene is the major product in this reaction.

Page : 320 , Block Name : Exercise

Q10.11 How will you bring about the following conversions?
(i) Ethanol to but-I-yne
(ii) Ethane to bromoethene
(iii) Propene to I-nitropropane
(iv) Toluene to benzyl alcohol
(v) Propene to propyne
(vi) Ethanol to ethyl fluoride
(vii) Bromomethane to propanone
(viii) But-I-ene to but-2-ene
(ix) I-Chlorobutane to n-octane
(x) Benzene to biphenyl

Answer.

(i)

$$\underset{\text{Ethanol}}{CH_3CH_2OH} \xrightarrow{SOCl_2,\ \text{Pyridine}} \underset{\text{Chloroethane}}{CH_3CH_2Cl} + SO_2 + HCl$$

$$\underset{\text{Ethyne}}{HC \equiv CH} + NaNH_2 \xrightarrow{\text{Liq } NH_3} \underset{\text{Sodium acetylide}}{HC \equiv C^{-}\,Na^{+}}$$

$$\underset{\text{Chloroethane}}{CH_3CH_2 - Cl} + HC \equiv C^{-}\,Na^{+} \longrightarrow \underset{\text{But}-1-\text{yne}}{CH_3CH_2C \equiv CH} + NaCl$$

(ii)



(iii)



(iv)



(v)



(vi)



(vii)



(viii)



(ix)



(x)



Page : 320 , Block Name : Exercise

Q10.12 Explain why
(i) the dipole moment of chlorobenzene is lower than that of cyclohexyl chloride?
(ii) alkyl halides, though polar, are immiscible with water?
(iii) Grignard reagents should be prepared under anhydrous conditions?

Answer. (i)



In chlorobenzene, the Cl-atom is linked to a sp^2 hybridized carbon atom. In cyclohexyl chloride, the Cl-atom is linked to a sp^3 hybridized carbon atom. Now, sp^2 hybridized carbon has more s-character than sp^2 hybridized carbon atom. Therefore, the former is more electronegative than the latter. Therefore, the density of electrons of C—CI bond near the Cl-atom is less in chlorobenzene than in cyclohexyl chloride.
Moreover, the —R effect of the benzene ring of chlorobenzene decreases the electron density of the C—CI bond near the Cl-atom. As a result, the polarity of the C—CI bond in chlorobenzene decreases. Hence, the dipole moment of chlorobenzene is lower than that of cyclohexyl chloride.
(ii) To be miscible with water, the solute-water force of attraction must be stronger than the solute-solute and water-water forces of attraction. Alkyl halides are polar molecules and so held together by dipole-dipole interactions. Similarly, strong H-bonds exist between the water

molecules. The new force of attraction between the alkyl halides and water molecules is weaker than the alkyl halide alkyl halide and water-water forces of attraction. Hence, alkyl halides (though polar) are immiscible with water.
(iii) Grignard reagents are very reactive. In the presence of moisture, they react to give alkanes.
$\mathrm{R\ MgX + H_2O \longrightarrow R - H + Mg(OH)X}$
Therefore, Grignard reagents should be prepared under anhydrous conditions.

Page : 320 , Block Name : Exercise

Q10.13 Give the uses of freon 12, DDT, carbon tetrachloride and iodoform.

Answer. Uses of Freon — 12
Freon-12 (dichlorodifluoromethane, $\mathrm{CF_2Cl_2}$) is commonly known as CFC. It is used as a refrigerant in refrigerators and air conditioners. It is also used in aerosol spray propellants such as body sprays, hair sprays, etc. However, it damages the ozone layer. Hence, its manufacture was banned in the United States and many other countries in 1994.
uses of DDT :
DDT (p, p dichlorodiphenyltrichloroethane) is one of the best known insecticides. It is very effective against mosquitoes and lice. But due its harmful effects, it was banned in the United States in 1973.

Uses of carbontetrachloride (CC14) :
(i) It is used for manufacturing refrigerants and propellants for aerosol cans.
(ii) It is used as feedstock in the synthesis of chlorofluorocarbons and other chemicals.
(iii) It is used as a solvent in the manufacture of pharmaceutical products.
(iv) Until the mid 1960's, carbon tetrachloride was widely used as a cleaning fluid, a degreasing agent in industries, a spot reamer in homes, and a fire extinguisher.

Uses Of iodoform (CHI,)
Iodoform was used earlier as an antiseptic, but now it has been replaced by other formulations-containing iodine-due to its objectionable smell. The antiseptic property of iodoform is only due to the liberation of free iodine when it comes in contact with the skin.

Page : 320 , Block Name : Exercise

Q10.14 Write the structure of the major organic product in each of the following reactions:

(i) $CH_3CH_2CH_2Cl + NaI \xrightarrow[\text{heat}]{\text{acetone}} \longrightarrow$

(ii) $(CH_3)_3CBr + KOH \xrightarrow[\text{heat}]{\text{ethanol}} \longrightarrow$

(iii) $CH_3CH(Br)CH_2CH_3 + NaOH \xrightarrow{\text{water}} \longrightarrow$

(iv) $CH_3CH_2Br + KCN \xrightarrow{\text{aq.ethanol}} \longrightarrow$

(v) $C_6H_5ONa + C_2H_5Cl \longrightarrow$

(vi) $CH_3CH_2CH_2OH + SOCl_2 \longrightarrow$

(vii) $CH_3CH_2CH = CH_2 + HBr \xrightarrow{\text{Peroxide}} \longrightarrow$

(viii) $CH_3CH = C(CH_3)_2 + HBr \longrightarrow$

Answer.

(i) $CH_3CH_2CH_2Cl + NaI \xrightarrow[\text{heat}]{\text{acetone}} CH_3CH_2CH_2I + NaCl$

1 – Chloropropane $\left(\begin{matrix}\text{Finkelstein}\\ \text{reaction}\end{matrix}\right)$ 1 – Iodopropane

(ii)

$(CH_3)_3CBr + KOH \xrightarrow[\text{heat}]{\text{ethanol}} CH_3 - \underset{\displaystyle CH_3}{\overset{\displaystyle |}{C}} = CH_2 + KBr + H_2O$

2-Bromo-2-methylpropane (Dehydrohalogenation)

2-Methylpropene

(iii) $CH_3CH(Br)CH_2CH_3 + NaOH \xrightarrow{\text{water}} CH_3CH(OH)CH_2CH_3 + NaBr$

2 – Bromobutane Butan – 2 – ol

(iv) $CH_3CH_2Br + KCN \xrightarrow[\text{(Nucleophilic substitution)}]{\text{aq. ethanol}} CH_3CH_2CN + KBr$

Bromobutane Cyanoethane

(v) $C_6H_5ONa + C_2H_5Cl \xrightarrow[\text{(Williamson synthesis)}]{} C_6H_5-O-C_2H_5 + NaCl$

Sodium phenoxide Chloroethane Phenetole

(vi) $CH_3CH_2CH_2OH + SOCl_2 \longrightarrow CH_3CH_2CH_2Cl + SO_2 + HCl$

1 – Propanol 1 – Chloropropane

(vii) $CH_3CH_2CH=CH_2 + HBr \xrightarrow[\text{(Anti-Markovnikov's addition)}]{\text{Peroxide}} CH_3CH_2CH_2CH_2-Br$

But – 1 – ene 1 – Bromobutane

(viii)

$CH_3CH=C(CH_3)_2 + HBr \xrightarrow[\text{addition)}]{\text{(Markovnikov}} CH_3-CH_2-\underset{\displaystyle CH_3}{\overset{\displaystyle Br}{\overset{|}{\underset{|}{C}}}}-CH_3$

2-Methylbut-2-ene 2-Bromo-2-methylbutane

Page : 320 , Block Name : Exercise

Q10.15 Write the mechanism of the following reaction:

$nBuBr + KCN \xrightarrow{EtOH-H_2O} nBuCN$

Answer. The given reaction is

$nBuBr + KCN \xrightarrow{EtOH-H_2O} nBuCN$

The given reaction is an S_N2 reaction. In this reaction, CN^- acts as the nucleophile and attacks the carbon atom to which ar is attached. CN^- ion is an ambident nucleophile and can attack through both C and N. In this case, it attacks through the C-atom.

$K^+CN^- + CH_3-CH_2-CH_2-\overset{\delta+}{CH_2}-\overset{\delta-}{Br} \xrightarrow[-Br^-]{} CH_3-CH_2-CH_2-CH_2-CN + KBr$

n-Butyl bromide *n*-Butyl cyanide

Page : 320 , Block Name : Exercise

Q10.16 Arrange the compounds of each set in order of reactivity towards SN2 displacement:
(i) 2-3romo-2-methylbutane, I-Bromopentane, 2-Bromopentane
(ii) I-Bromo-3-methylbutane, 2-3romo-2-methylbutane, 3-3romo-2- methylbutane
(iii) I-aromobutane, I-Bromo-2,2-dimethylpropane, I-Bromo-2-methylbutane, bromo-3-methylbutane.

Answer. (i)



An S_N2 reaction involves the approaching of the nucleophile to the carbon atom to which the leaving group is attached. When the nucleophile is sterically hindered, then the reactivity towards S_N2 displacement decreases. Due to the presence of substituents, hindrance to the approaching nucleophile increases in the following order.

I-Bromopentane < 2-bromopentane < 2-3romo-2-methylbutane

Hence, the increasing order of reactivity towards S_N2 displacement is:

2-3romo-2-methylbutane < 2-Bromopentane < I-Bromopentane

(ii)



Since steric hindrance in alkyl halides increases in the order of $1^0 < 2^0 < 3^0$, the increasing order of reactivity towards SN2 displacement is

$3^0 < 2^0 < 1^0$

Hence, the given set of compounds can be arranged in the increasing order of their reactivity towards S_N^2 displacement as:

2-aromo-2-methylbutane < 2-Bromo-3-methylbutane < I-aromo-3-methylbutane

[2-3romo-3-methylbutane is incorrectly given in NCERT]

(iii)



The steric hindrance to the nucleophile in the S_N2 mechanism increases with a decrease in the distance of the substituents from the atom containing the leaving group. Further, the steric hindrance increases with an increase in the number of substituents. Therefore, the increasing order of steric hindrances in the given compounds is as below:

I-Bromobutane < I-Bromo-3-methylbutane < I-Bromo-2-methylbutane

< I-Bromo-2, 2-dimethylpropane

Hence, the increasing order of reactivity of the given compounds towards SN2 displacement is: I-Bromo-2, 2-dimethylpropane < I-Bromo-2-methylbutane < I-aromo-3 methylbutane < I-Bromobutane

Page : 320 , Block Name : Exercise

Q10.17 Out of $C_6H_5CH_2Cl$ and $C_6H_5CHClC_6H_5$, which is more easily hydrolysed by aqueous KOH?

Answer.



Hydrolysis by aqueous KOH proceeds through the formation of carbocation. If carbocation is stable, then the compound is easily hydrolyzed by aqueous KOH. Now, $C_6H_5CH_2Cl$ forms 1° -carbocation While $C_6H_5CHCLC_6H_5$ forms 2° -carbocation, which is more stable than 2°-carbocation. Hence, $C_6H_5CHCIC_6H_5$ 6 is hydrolyzed more easily than $C_eH_3CH_2Cl$ by aqueous KOH.

Page : 320 , Block Name : Exercise

Q10.18 p-Dichlorobenzene has higher m.p. and lower solubility than those of o- and m-isomers. Discuss.

Answer.



p-Dichlorobenzene is more symmetrical than 0-and m-isomers. For this reason, it fits more closely than O_and m_isomers in the crystal lattice. Therefore, more energy is required to break the crystal lattice of p-dichlorobenzene. As a result, p-dichlorobenzene has a higher melting point and lower solubility than 0-and m-isomers.

Page : 320 , Block Name : Exercise

Q10.19 How the following conversions can be carried out?
(i) Propene to propan-l-ol

(ii) Ethanol to but-I-yne
(iii) I-Bromopropane to 2-bromopropane
(iv) Toluene to benzyl alcohol
(v) Benzene to 4-bromonitrobenzene
(vi) Benzyl alcohol to 2-phenylethanoic acid
(vii) Ethanol to propanenitrile
(viii) Aniline to chlorobenzene
(ix) 2-Chlorobutane to 3, 4-dimethylhexane
(x) 2-Methyl-1-propene to 2-chloro-2-methylpropane
(xi) Ethyl chloride to propanoic acid
(xii) But-I-ene to n-butyliodide
(xiii) 2-Chloropropane to I-propanol
(xiv) Isopropyl alcohol to iodoform
(xv) Chlorobenzene to p-nitrophenol
(xvi) 2-3romopropane to I-bromopropane
(xvii) Chloroethane to butane
(xviii) Benzene to diphenyl
(xix) tert-Butyl bromide to isobutyl bromide
(xx) Aniline to phenylisocyanide

Answer.

(i)



(ii)


CH3CH2OH
red P / Br2
CH3CH2Br
Mg / ether
CH3CH2MgBr
Ethanol
Ethyl bromide
Ethyl magnesium bromide
ether
(Grignard reaction)
H+ / H2O
CH3CH2CH2CH2OH
[CH3CH2CH2CH2OMgBr]
1 – Butanol
Addition product
red P / I2
CH3CH2CH2CH2 — I
KOH (alc) / Δ
(Dehydrohalogenation)
CH3CH2CH = CH2
Iodobutane
But – 1 – ene
(Electrophilic addition)
Br4 / CCl4
CH3CH2CH — CH2
Br Br
NaNH2 , liq NH3
196 K
(Dehydrohalogenation)
CH3CH2CH ≡ CH
But – 1 – yne
1, 2 – Dibromobutane

(iii)


CH3 — CH2 — CH2 — Br
1 – Bromopropane
KOH (alc) / Δ
(Dehydrohalogenation)
CH3 — CH = CH2
Propene
HBr
Markovnikov addition
Br
CH3 — CH — CH3
2 – bromopropane

(iv)


CH3
Cl2 / UV light
or heat
CH2Cl
Aq. KOH / Δ
(nucleophilic substitution)
CH2OH
Toluene
Benzyl chloride
Benzyl alcohol

(v)


Br2 / FeBr3
Dark
HNO3 / H2SO4
(Nitration)
Br
Br
NO2
Benzene
Bromobenzene
4 – Bromonitrobenzene

(vi)



(vii)



(viii)



(ix)



(x)



(xi)


CH3 — CH2 — Cl
Ethylchloride
KCN, aq. ethanol
(nucleophilic substitution)
CH3 — CH2 — CN
Propanenitrile
+ KCl
(Hydrolysis)
H+ / H2O
CH3 — CH2 — COOH
Propanoic acid

(xii)


CH3 — CH2 — CH = CH2
But – 1 – ene
HBr / Peroxide
(Anti - Markovnikov addition)
CH3 — CH2 — CH2 — CH2 — Br
1 – Bromobutane
Finkelstein reaction
NaI, dry acetone
CH3 — CH2 — CH2 — CH2 — I
n – Butyliodide

(xiii)


Cl
CH3 — CH — CH3
2 – Chloropropane
KOH (alc) / Δ
(Dehydrohalogenation)
CH3 — CH = CH2
Propene
+ HCl
HBr / Peroxide
(Anti - Markovnikov addition)
CH3 — CH2 — CH2 — Br
1 – Bromopropane
Aq. KOH / Δ
(Nucleophilic substitution)
CH3 — CH2 — CH2 — OH
1 – Propanol

(xiv)


OH
CH3 — CH — CH3
CrO3
(Oxidation)
O
CH3 — C — CH3
Isopropyl alcohol
Propanone
NaOI
(Iodoform reaction)
O
CH3 — C — ONa
+
CHI3
Iodoform

(xv)


Cl
HNO3 / H2SO4
(Nitration)
Chlorobenzene
NO2
p – Chloronitrobenzene
(Major product)
m – Chloronitrobenzene
o – Chloronitrobenzene


Cl
(i) NaOH, 433 K
(ii) dil HCl
OH
NO2
p – Chloronitrobenzene
p – Nitrophenol

(xvi)


Br
CH3 — CH — CH3
2 – Bromopropane
KOH (alc) / Δ
Dehydrohalogenation
CH3 — CH = CH2 + HBr
Propene
(Anti–Markovnikov addition)
HBr / Peroxide
CH3 — CH2 — CH2 — Br
1 – Bromopropane

(xvii)



(xviii)



(xix)



(xx)



Page : 320 , Block Name : Exercise

Q10.20 The treatment of alkyl chlorides with aqueous KOH leads to the formation of alcohols but in the presence of alcoholic KOH, alkenes are major products. Explain.

Answer. In an aqueous solution, KOH almost completely ionizes to give OH^- ions. OH^- ion is a strong nucleophile, which leads the alkyl chloride to undergo a substitution reaction to form alcohol.

$$R - Cl + KOH_{(aq)} \longrightarrow R - OH + KCl$$

On the other hand, an alcoholic solution of KOH contains alkoxide RO^- ion, which is a strong base. Thus, it can abstract a hydrogen from the ß-carbon of the alkyl chloride and form an alkene by eliminating a molecule of HCI.

$$R - CH_2 - CH_2 - Cl + KOH(alc) \longrightarrow R - CH = CH_2 + KCl + H_2O$$

OH^- ion is a much weaker base than RO^- ion. Also, OH^- ion is highly solvated in an aqueous solution and as a result, the basic character of ion decreases. Therefore, it cannot abstract a hydrogen from the ß-carbon.

Page : 321 , Block Name : Exercise

Q10.21 Primary alkyl halide C_4H_9Br (a) reacted with alcoholic KOH to give compound (b).Compound (b) is reacted with Har to give (c) which is an isomer of (a). When (a) is reacted with sodium metal it gives compound (d), C_8H_{18} which is different from the compound formed when n-

butyl bromide is reacted with sodium. Give the structural formula of (a) and write the equations for all the reactions.

Answer. There are two primary alkyl halides having the formula, C4H9Br. They are n —bulyl bromide and isobutyl bromide.



Therefore, compound (a) is either II—butyl bromide or isobutyl bromide.
Now, compound (a) reacts with Na metal to give compound (b) of molecular formula, C_8H_{18} which is different from the compound formed when n—butyl bromide reacts with Na metal. Hence. compound (a) must be isobutvl bromide.





Thus, compound (d) is 2, 5–dimethylhexane.

It is given that compound (a) reacts with alcoholic KOH to give compound (b). Hence, compound (b) is 2–methylpropene.



Also, compound (b) reacts with HBr to give compound (c) which is an isomer of (a). Hence, compound (c) is 2–bromo–2–methylpropane.



Page : 321 , Block Name : Exercise

Q10.22 What happens when

(i) n-butyl chloride is treated with alcoholic KOH,
(ii) bromobenzene is treated with Mg in the presence of dry ether,
(iii) chlorobenzene is subjected to hydrolysis,
(iv) ethyl chloride is treated with aqueous KOH,
(v) methyl bromide is treated with sodium in the presence of dry ether,
(vi) methyl chloride is treated with KCN.

Answer.

(i) When n–butyl chloride is treated with alcoholic KOH, the formation of but–l–ene takes place. This reaction is a dehydrohalogenation reaction.

$$CH_3-CH_2-CH_2-CH_2-Cl \xrightarrow[(\text{Dehydrohalogenation})]{KOH(alc)/\Delta} CH_3-CH_2-CH=CH_2 + KCl + H_2O$$

n – Butyl chloride — But-l-ene

(ii) When bromobenzene is treated with Mg in the presence of dry ether, phenylmagnesium bromide is formed.



Bromobenzene — Phenylmagensium bromide

(iii) Chlorobenzene does not undergo hydrolysis under normal conditions. However, it undergoes hydrolysis when heated in an aqueous sodium hydroxide solution at a temperature of 623 K and a pressure of 300 atm to form phenol.



Chlorobenzene — Phenol

(iv) When ethyl chloride is treated with aqueous KOH, it undergoes hydrolysis to form ethanol.

$$CH_3-CH_2-Cl \xrightarrow[(\text{Hydrolysis})]{KOH_{(aq)}} CH_3-CH_2-OH + KCl$$

Ethyl chloride — Ethanol

(v) When methyl bromide is treated with sodium in the presence of dry ether, ethane is formed. This reaction is known as the Wurtz reaction.

$$2\,CH_3-Br + 2\,Na \xrightarrow[(\text{Wurtz reaction})]{\text{Dry ether}} CH_3-CH_3 + 2NaBr$$

Methyl bromide — Ethane

(vi) When methyl chloride is treated with KCN, it undergoes a substitution reaction to give methyl cyanide.

$$CH_3-Cl + KCN \xrightarrow[\text{Nucleophilic substitution}]{} CH_3-CN + KCl$$

Methyl chloride — Methyl cyanide