NCERT SOLUTIONS

CLASS - 12th





aglasem.com


Class : 12th
Subject : Chemistry
Chapter : 11
Chapter Name : Alcohols, Phenols Phenols and Ethers

Q11.1 Classify the following as primary, secondary and tertiary alcohols:


(i)




(ii) 

(iii) 


(iv)



(v)



(vi)



Answer. Primary alcohol (i), (ii), (iii)
Secondary alcohol (iv), (v)
Tertiary alcohol (vi)

Page : 325 , Block Name : Intext Questions


Q11.2 Identify allylic alcohols in the above question.

Answer. The alcohols given in (ii) and (vi) are allylic alcohols.


Page : 325 , Block Name : Intext Questions

Q11.3 Name the following compounds according to IUPAC system.


(i)




(ii)




(iii)



(iv)



(v)




Answer. (i) 3-Chloromethyl-2-isopropylpentan-1-ol
(ii) 2, 5-Dimethylhexane-1, 3-diol
(iii) 3-Bromocyclohexanol
(iv) Hex-1-en-3-ol
(v) 2-Bromo-3-methylbut-2-en-1-ol


Page : 328 , Block Name : Intext Questions

Q11.4 Show how are the following alcohols prepared by the reaction of a suitable Grignard reagent on methanal?

(i)




(ii)



Answer.

(i)




(ii)




Page : 328 , Block Name : Intext Questions

Q11.5 Write structures of the products of the following reactions:

(i) $\mathrm{CH_3 - CH = CH_2} \overset{\mathrm{H_2O/H^+}}{\rightarrow}$

(ii) 

(iii) 

Answer. Missing

Page : 328 , Block Name : Intext Questions

Q11.6 Give structures of the products you would expect when each of the following alcohol reacts with (a) HCl – $ZnCl_2$ (b) HBr and (c) $SOCl_2$.


(i) Butan-1-ol

(ii) 2-Methylbutan-2-ol

Answer.


(a)

(i)



Primary alcohols do not react appreciably with Lucas' reagent ($HCl\text{-}ZnCl_2$) at room temperature.

(ii)




2 – Methylbutan – 2 – ol (3°) — 2 – Chloro – 2 – Methylbutane (White turbidity)


Tertiary alcohols react immediately with Lucas' reagent.

(b)

(i)




(ii)




2 – Methylbutan – 2 – ol (3°) — 2 – Bromo – 2 – Methylbutane

(c)

(i)



(ii)




2 – Methylbutan – 2 – ol — 2 – Chloro – 2 – Methylbutane

Page : 343 , Block Name : Intext Questions

Q11.7 Predict the major product of acid catalysed dehydration of
(i) 1-methylcyclohexanol and (ii) butan-1-ol

Answer.

i.



1 – Methylcyclohexanol 1 – Methylcyclohexene (Major product)

(ii)


$$CH_3CH_2CH_2CH_2OH \xrightarrow[H^+]{\text{Dehydration}} CH_3CH=CHCH_3 + H_2O$$

Butan-1-ol But-2-ene (Major product)


Page : 343 , Block Name : Intext Questions

Q11.8 Ortho and para nitrophenols are more acidic than phenol. Draw the resonance structures of the corresponding phenoxide ions.


Answer.



Resonance structure of the phenoxide ion



Resonance structures of *p*-nitrophenoxide ion




Resonance structures of m-nitrophenoxide ion. It can be observed that the presence of nitro groups increases the stability of phenoxide Ion.

Page : 343 , Block Name : Intext Questions


Q11.9 Write the equations involved in the following reactions:
(i) Reimer - Tiemann reaction
(ii) Kolbe's reaction

Answer.

i. **Reimer-Tiemann reaction**




ii. **Kolbe's reaction**




Page : 343 , Block Name : Intext Questions


Q11.10 Write the reactions of Williamson synthesis of 2-ethoxy-3-methylpentane starting from ethanol and 3-methylpentan-2-ol.

Answer. In Williamson synthesis, an alkyl halide reacts with an alkoxide ion. Also, it is an SN2 reaction. In the reaction, alkyl halides should be primary having the least steric hindrance. Hence, an alkyl halide is obtained from ethanol and alkoxide ion from 3- methylpentan-2-ol.





3 – Methylpentan – 2 – ol Sodium 3 – Methylpentan – 2 – oxide




2 – Ethoxy – 3 – methylpentane


Page : 350 , Block Name : Intext Questions

Q11.11 Which of the following is an appropriate set of reactants for the preparation of 1-methoxy-4-nitrobenzene and why?


(i)



(ii)



Answer. Set (ii) is an appropriate set of reactants for the preparation of I-methoxy-4- nitrobenzene.




1 – Methoxy – 4 – nitrobenzene


In set (i), sodium methoxide (CH30Na) is a strong nucleophile as well as a strong base.
Hence, an elimination reaction predominates over a substitution reaction.

Page : 350 , Block Name : Intext Questions

Q11.12 Predict the products of the following reactions:

(i) $CH_3-CH_2-CH_2-O-CH_3+HBr \rightarrow$

(ii) 

(iii) 

(iv) $(CH_3)_3C-OC_2H_5 \xrightarrow{HI}$

Answer.

Answer

(i)


$CH_3-CH_2-CH_2-O-CH_3 + HBr \longrightarrow$

n-propylmethyl ether

$CH_3-CH_2-CH_2-OH + CH_3-Br$


Propanol Bromomethane

(ii)



Ethoxybenzene Phenol Bromoethane

(iii)



Ethoxybenzene 4 – Ethoxynitrobenzene (Major) 2 – Ethoxynitrobenzene (Minor)

(iv)


$(CH_3)_3C-OC_2H_5 \xrightarrow{HI} (CH_3)_3C-I + C_2H_5OH$

tert-Butyl ethyl ether *tert*-Butyliodide Ethanol


Page : 350 , Block Name : Intext Questions

Q11.1 Write IUPAC names of the following compounds:


(i)




(ii)




(iii)



(iv)




(v)




(vi)







(vii)



(viii)




(ix)



(x) $C_6H_5-O-C_2H_5$

(xi) $C_6H_5-O-C_7H_{15}(n-)$

(xii)



Answer. (i) 2, 2, 4-Trimethylpentan-3-ol
(ii) 5-Ethylheptane-2, 4-diol
(iii) Butane-2, 3-diol
(iv) Propane-I, 2, 3-triol
(v) 2-Methylphenol
(vi) 4-Methylphenol
(vii) 2, 5-Dimethylphenol
(viii) 2, 6-Dimethylphenol
(ix) I-Methoxy-2-methylpropane
(x) Ethoxybenzene
(xi) I-Phenylheptane
(xii) 2-Ethoxybutane


Page : 352 , Block Name : Exercise

Q11.2 Write structures of the compounds whose IUPAC names are as follows:
(i) 2-Methylbutan-2-ol
(ii) 1-Phenylpropan-2-ol
(iii) 3,5-Dimethylhexane –1, 3, 5-triol
(iv) 2,3 – Diethylphenol
(v) 1 – Ethoxypropane
(vi) 2-Ethoxy-3-methylpentane
(vii) Cyclohexylmethanol


(viii) 3-Cyclohexylpentan-3-ol
(ix) Cyclopent-3-en-1-ol
(x) 4-Chloro-3-ethylbutan-1-ol

Answer.


(i)




(ii)




(iii)




(iv)




(v)




(vi)



(vii)



(viii)




(ix)





(x)



Page : 352 , Block Name : Exercise

Q11.3 (i) Draw the structures of all isomeric alcohols of molecular formula C5H12O and give their IUPAC names.
ii) Classify the isomers of alcohols in question 11.3 (i) as primary, secondary and tertiary alcohols.


Answer.

(i) The structures of all isomeric alcohols of molecular formula, $C_5H_{12}O$ are shown below:

(a) $CH_3-CH_2-CH_2-CH_2-CH_2-OH$


Pentan-1-ol (1°)

(b)




2-Methylbutan-1-ol (1°)

(c)




3-Methylbutan-1-ol (1°)

(d)




2, 2-Dimethylpropan-1-ol (1°)

(e)




Pentan-2-ol (2°)

(f)



3-Methylbutan-2-ol (2°)


(g)





Pentan-3-ol (2°)

(h)



2-Methylbutan-2-ol (3°)

(ii) Primary alcohol: Pentan-1-ol; 2-Methylbutan-1-ol;

3-Methylbutan-1-ol; 2, 2−Dimethylpropan-1-ol

Secondary alcohol: Pentan-2-ol; 3-Methylbutan-2-ol;


Pentan-3-ol

Tertiary alcohol: 2-methylbutan-2-ol

Page : 352 , Block Name :Exercise

Q11.4 Explain why propanol has higher boiling point than that of the hydrocarbon, butane?

Answer. Propanol undergoes intermolecular H-bonding because of the presence of —OH group.
On the other hand, butane does not




Therefore, extra energy is required to break hydrogen bonds. For this reason, propanol has a higher boiling point than hydrocarbon butane.

Page : 352 , Block Name :Exercise

Q11.5 Alcohols are comparatively more soluble in water than hydrocarbons of comparable molecular masses. Explain this fact.

Answer. Alcohols form H-bonds with water due to the presence of hydrocarbons cannot form H-bonds with water. — OH group. However,




As a result, alcohols are comparatively more soluble in water than hydrocarbons of comparable molecular masses.

Page : 352 , Block Name : Exercise

Q11.6 What is meant by hydroboration-oxidation reaction? Illustrate it with an example.


Answer. The addition Of borane followed by Oxidation is known as the hydroboration-oxidation reaction.
For example, propan-l-ol is produced by the hydroboration-oxidation reaction
of propene. In this reaction, propene reacts with diborane (BH3)2 to form trialkyl borane as an addition product.
This addition product is oxidized to alcohol by hydrogen peroxide in the presence of aqueous sodium hydroxide.



Page : 352 , Block Name : Exercise

Q11.7 Give the structures and IUPAC names of monohydric phenols of molecular formula, C_7H_aO


Answer.



Page : 352 , Block Name : Exercise

Q11.8 While separating a mixture of ortho and para nitrophenols by steam distillation, name the isomer which will be steam volatile. Give reason.


Answer. Intramolecular H-bonding is present in 0-nitrophenol and p-nitrophenol. In p- nitrophenol, the molecules are strongly associated due to the presence Of intermolecular bonding. Hence, 0-nitrophenol is steam volatile.




Page : 352 , Block Name : Exercise

Q11.9 Give the equations of reactions for the preparation of phenol from cumene.

Answer. To prepare phenol, cumene is first oxidized in the presence of air of cumene hydroperoxide.




Then, cumene hydroxide is treated with dilute acid to prepare phenol and acetone as by- products.



Page : 352 , Block Name : Exercise

Q11.10 Write chemical reaction for the preparation of phenol from chlorobenzene

Answer. Chlorobenzene is fused with NaOH (at 623 K and 320 atm pressure) to produce sodium phenoxide, which gives phenol on acidification.



Page : 352 , Block Name : Exercise


Q11.11 Write the mechanism of hydration of ethene to yield ethanol.

Answer.

The mechanism of hydration of ethene to form ethanol involves three steps.


Step 1:

Protonation of ethene to form carbocation by electrophilic attack of H_3O^+:




Step 2:

Nucleophilic attack of water on carbocation:



Step 3:


Deprotonation to form ethanol:



Page : 352 , Block Name : Exercise

Q11.12 You are given benzene, conc. H2SO4 and NaOH. Write the equations for the preparation of phenol using these reagents.

Answer.




Page : 353 , Block Name : Exercise

Q11.13 Show how will you synthesise:
(i) 1-phenylethanol from a suitable alkene.
(ii) cyclohexylmethanol using an alkyl halide by an SN2 reaction.
(iii) pentan-1-ol using a suitable alkyl halide?


Answer.

(i) By acid-catalyzed hydration of ethylbenzene (styrene), 1-phenylethanol can be synthesized.



Phenylethene 1 – phenylethanol

(ii) When chloromethylcyclohexane is treated with sodium hydroxide, cyclohexylmethanol is obtained.



Chloromethylcyclohexane Cyclohexylmethanol

(iii) When 1-chloropentane is treated with NaOH, pentan-1-ol is produced.

$$CH_3CH_2CH_2CH_2CH_2Cl + NaOH \longrightarrow CH_3CH_2CH_2CH_2CH_2OH + NaCl$$

1-Chloropentane Pantan-1-ol


Page : 353 , Block Name : Exercise

Q11.14 Give two reactions that show the acidic nature of phenol. Compare acidity of phenol with that of ethanol.

Answer.


The acidic nature of phenol can be represented by the following two reactions:

(i) Phenol reacts with sodium to give sodium phenoxide, liberating H_2.




Phenol Sodium phenoxide

(ii) Phenol reacts with sodium hydroxide to give sodium phenoxide and water as by-products.



Sodium phenoxide


The acidity of phenol is more than that of ethanol. This is because after losing a proton, the phenoxide ion undergoes resonance and gets stabilized whereas ethoxide ion does not.



Page : 353 , Block Name : Exercise

Q11.15 Explain why is ortho nitrophenol more acidic than ortho methoxyphenol ?

Answer.



o – Nitrophenol o – Methoxyphenol


The nitro-group is an electron-withdrawing group. The presence Of this group in the ortho position decreases the electron density in the bond. As a result, it is easier to lose a proton. Also, the 0-nitrophenoxide ion formed after the loss of protons is stabilized by resonance. Hence, ortho nitrophenol is a stronger acid.
On the other hand, methoxy group is an electron-releasing group. Thus, it increases the electron density in the bond and hence, the proton cannot be given out easily.
For this reason, ortho-nitrophenol is more acidic than ortho-methoxyphenol.

Page : 353 , Block Name : Exercise

Q11.16 Explain how does the –OH group attached to a carbon of benzene ring activate it towards electrophilic substitution?

Answer. The —OH group is an electron-donating group. Thus, it increases the electron density in the benzene ring

as shown in the given resonance structure of phenol.




As a result, the benzene ring is activated towards electrophilic substitution.

Page : 353 , Block Name : Exercise

Q11.17 Give equations of the following reactions:
(i) Oxidation of propan-1-ol with alkaline KMnO4 solution.
(ii) Bromine in CS2 with phenol.
(iii) Dilute HNO3 with phenol.
(iv) Treating phenol with chloroform in presence of aqueous NaOH.


Answer.




(i)

(ii)

(iii)




(iv)




Page : 353 , Block Name : Exercise

Q11.18 Explain the following with an example.
(i) Kolbe’s reaction.
(ii) Reimer-Tiemann reaction.
(iii) Williamson ether synthesis.
(iv) Unsymmetrical ether

Answer. (i) Kolbe's reaction:
When phenol is treated with sodium hydroxide, sodium phenoxide is produced. This sodium phenoxide when treated with carbon dioxide, followed by acidification, undergoes electrophilic substitution to give ortho-hydroxybenzoic acid as the main product. This reaction is known as Kolbe's reaction.




(ii) Reimer—Tiemann reaction:
When phenol is treated with chloroform (CHCh) in the presence of sodium hydroxide, a —CHO group is introduced at the ortho position of the benzene ring.




This reaction is known as the Reimer-Tiemann reaction.

The intermediate is hydrolyzed in the presence of alkalis to produce salicyclaldehyde.



(iii) Williamson ether synthesis:

Williamson ether synthesis is a laboratory method to prepare symmetrical and unsymmetrical ethers by allowing alkyl halides to react with sodium alkoxides.



This reaction involves S_N2 attack of the alkoxide ion on the alkyl halide. Better results are obtained in case of primary alkyl halides.



If the alkyl halide is secondary or tertiary, then elimination competes over substitution.

(iv) Unsymmetrical ether:

An unsymmetrical ether is an ether where two groups on the two sides of an oxygen atom differ (i.e., have an unequal number of carbon atoms). For example: ethyl methyl ether ($CH_3–O–CH_2CH_3$).

Page : 353 , Block Name : Exercise


Q11.19 Write the mechanism of acid dehydration of ethanol to yield ethene

Answer.

The mechanism of acid dehydration of ethanol to yield ethene involves the following three steps:


Step 1:

Protonation of ethanol to form ethyl oxonium ion:




Step 2:

Formation of carbocation (rate determining step):



Step 3:

Elimination of a proton to form ethene:



The acid consumed in step 1 is released in Step 3. After the formation of ethene, it is removed to shift the equilibrium in a forward direction.


Page : 353 , Block Name : Exercise

Q11.20 How are the following conversions carried out?
(i) Propene → Propan-2-ol.
(ii) Benzyl chloride → Benzyl alcohol.
(iii) Ethyl magnesium chloride → Propan-1-ol.
(iv) Methyl magnesium bromide → 2-Methylpropan-2-ol

Answer.


Answer

(i) If propene is allowed to react with water in the presence of an acid as a catalyst, then propan-2-ol is obtained.




Propene — Propan – 2 – ol

(ii) If benzyl chloride is treated with NaOH (followed by acidification) then benzyl alcohol is produced.



Benzyl chloride — Benzyl alcohol

(iii) When ethyl magnesium chloride is treated with methanal, an adduct is the produced which gives propan-1-ol on hydrolysis.




Adduct

$Mg(OH)Cl + C_3H_7 - OH$

Propan – 1 – ol

(iv) When methyl magnesium bromide is treated with propane, an adduct is the product which gives 2-methylpropane-2-ol on hydrolysis.



Adduct

2 – Methylpropan – 2 – ol

Page : 353 , Block Name : Exercise

Q11.21 Name the reagents used in the following reactions:
(i) Oxidation of a primary alcohol to carboxylic acid.
(ii) Oxidation of a primary alcohol to aldehyde.


(iii) Bromination of phenol to 2,4,6-tribromophenol.
(iv) Benzyl alcohol to benzoic acid.
(v) Dehydration of propan-2-ol to propene.
(vi) Butan-2-one to butan-2-ol.

Answer. (i) Acidified potassium permanganate
(ii) Pyridinium chlorochromate (PCC)
(iii) Bromine water
(iv) Acidified potassium permanganate
(v) 85% phosphoric acid
(vi) NaBH4 or LiAlH4

Page : 353 , Block Name : Exercise


Q11.22 Give reason for the higher boiling point of ethanol in comparison to methoxymethane.

Answer. Ethanol undergoes intermolecular H-bonding due to the presence of —OH group, resulting in the association Of molecules. Extra energy is required to break these hydrogen bonds. On the other hand, methoxymethane does not undergo H-bonding.
Hence, the boiling point of ethanol is higher than that of methoxymethane.



Page : 354 , Block Name : Exercise

Q11.23 Give IUPAC names of the following ethers




Answer. (i) I-Ethoxy-2-methylpropane
(ii) 2-chloro-l-methoxyethane
(iii) 4-Nitroanisole
(iv) I-Methoxypropane
(v) 1-Ethoxy-4, 4-dimethylcyclohexane
(vi) Ethoxybenzene

Page : 354 , Block Name : Exercise


Q11.24 Write the names of reagents and equations for the preparation of the following ethers by Williamson's synthesis:
(i) 1-Propoxypropane
(ii) Ethoxybenzene
(iii) 2-Methoxy-2-methylpropane
(iv) 1-Methoxyethane

Answer.


(i)




(ii)



(iii)




(iv)




Page : 354 , Block Name : Exercise

Q11.25 Illustrate with examples the limitations of Williamson synthesis for the preparation of certain types of ethers.

Answer. The reaction of Williamson synthesis involves SN2 attack of an alkoxide ion on a primary alkyl halide.



But if secondary or tertiary alkyl halides are taken in place of primary alkyl halides, then elimination would compete over substitution. As a result, alkenes would be produced. This is because alkoxides are nucleophiles as well as strong bases. Hence, they react with alkyl halides, which results in an elimination reaction.




Page : 354 , Block Name : Exercise

Q11.26 How is 1-propoxypropane synthesised from propan-1-ol? Write mechanism of this reaction

Answer.

1-propoxypropane can be synthesized from propan-1-ol by dehydration.


Propan-1-ol undergoes dehydration in the presence of protic acids (such as H_2SO_4, H_3PO_4) to give 1-propoxypropane.



Propane-1-ol 1-Propoxypropane


The mechanism of this reaction involves the following three steps:

Step 1: Protonation




Popan – 1 – ol

Step 2: Nucleophilic attack



Step 3: Deprotonation



1 – Propoxypropane

Page : 354 , Block Name : Exercise

Q11.27 Preparation of ethers by acid dehydration of secondary or tertiary alcohols is not a suitable method. Give reason.


Answer. The formation of ethers by dehydration of alcohol is a bimolecular reaction (SN2) involving the attack of an alcohol molecule on a protonated alcohol molecule. In the method, the alkyl group should be unhindered. In case of secondary or tertiary alcohols, the alkyl group is hindered. As a result, elimination dominates substitution.
Hence, in place of ethers, alkenes are formed.

Page : 354 , Block Name : Exercise


Q11.28 Write the equation of the reaction of hydrogen iodide with:
(i) 1-propoxypropane (ii) methoxybenzene and (iii) benzyl ethyl ether

Answer.


(i)



(ii)



(iii)




Page : 354 , Block Name : Exercise


Q11.29 Explain the fact that in aryl alkyl ethers
(i) the alkoxy group activates the benzene ring towards electrophilic substitution and
(ii) it directs the incoming substituents to ortho and para positions in benzene ring.

Answer.

(i)



In aryl alkyl ethers, due to the +R effect of the alkoxy group, the electron density in the benzene ring increases as shown in the following resonance structure.



Thus, benzene is activated towards electrophilic substitution by the alkoxy group.

(ii) It can also be observed from the resonance structures that the electron density increases more at the ortho and para positions than at the meta position. As a result, the incoming substituents are directed to the ortho and para positions in the benzene.

Page : 354 , Block Name : Exercise

Q11.30 Write the mechanism of the reaction of HI with methoxymethane.


Answer.

The mechanism of the reaction of HI with methoxymethane involves the following steps:

Step1: Protonation of methoxymethane:




Step2: Nucleophilic attack of I^{-}:



Iodomethane methanol

Step3:

When HI is in excess and the reaction is carried out at a high temperature, the methanol formed in the second step reacts with another HI molecule and gets converted to methyl iodide






Page : 354 , Block Name : Exercise


Q11.31 Write equations of the following reactions:
(i) Friedel-Crafts reaction – alkylation of anisole.
(ii) Nitration of anisole.
(iii) Bromination of anisole in ethanoic acid medium.
(iv) Friedel-Crafts acetylation of anisole.

Answer.


(i)




(ii)



(iii)




(iv)




Page : 354 , Block Name : Exercise

Q11.32 Show how would you synthesise the following alcohols from appropriate alkenes?


(i)




(ii)



(iii)




(iv)




Answer.

The given alcohols can be synthesized by applying Markovnikov's rule of acid-catalyzed hydration of appropriate alkenes.


(i)




(ii)



(iii)




Acid-catalyzed hydration of pent-2-ene also produces pentan-2-ol but along with pentan-3-ol.




Thus, the first reaction is preferred over the second one to get pentan-2-ol.

(iv)



Page : 354 , Block Name : Exercise

Q11.33 When 3-methylbutan-2-ol is treated with HBr, the following reaction takes place:



Give a mechanism for this reaction.
(Hint : The secondary carbocation formed in step II rearranges to a more stable tertiary carbocation by a hydride ion shift from 3rd carbon atom.)

Answer.


The mechanism of the given reaction involves the following steps:

Step 1: Protonation




3 – Methylbutan – 2 – ol

Step 2: Formation of 2° carbocation by the elimination of a water molecule




2° Carbocation

Step 3: Re-arrangement by the hydride-ion shift



(less stable) 3° Carbocation (more stable)

Step 4: Nucleophilic attack



2 – Bromo – 2 – methylbutane

Page : 354 , Block Name : Exercise