SECTION-A

Q1. $x^2 \frac{d^2y}{dx^2} = \left\{1 + \left(\frac{dy}{dx}\right)^2\right\}^4$

ORDER = 2

DEGREE = 1

Q2. $f(x) = x+7 \; ; \; g(x) = x-7$

$$fog(x) = f(g(x))$$
$$= f(x-7)$$
$$= (x-7)+7$$
$$= x \qquad \forall \; x \in R$$

$$\frac{d}{dx} fog(x) = \frac{d}{dx}(x) = 1$$

$$\boxed{\frac{d}{dx} fog(x) = 1}$$

Q3 $2\begin{bmatrix} 1 & 3 \\ 0 & x \end{bmatrix} + \begin{bmatrix} y & 0 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 6 \\ 1 & 8 \end{bmatrix}$

$\Rightarrow \begin{bmatrix} 2 & 6 \\ 0 & 2x \end{bmatrix} + \begin{bmatrix} y & 0 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 6 \\ 1 & 8 \end{bmatrix}$

$\Rightarrow \begin{bmatrix} 2+y & 6 \\ 1 & 2x+2 \end{bmatrix} = \begin{bmatrix} 5 & 6 \\ 1 & 8 \end{bmatrix}$

comparing corresponding elements of each matrix,

$2+y = 5$ $\qquad$ $2x+2 = 8$

$\boxed{y = 3}$ $\qquad$ $\boxed{x = 3}$

$x - y = 3 - 3$

$\boxed{x - y = 0}$

Q4 $\vec{r} = \vec{a} + \lambda\vec{b}$

$\boxed{\vec{r} = (3\hat{i} + 4\hat{j} + 5\hat{k}) + \lambda(2\hat{i} + 2\hat{j} - 3\hat{k})}$

where $\vec{a}$ = position vector of lot or line

$\vec{b}$ = parallel vector to line

Also $\vec{r} = (3+2\lambda)\hat{i} + (4+2\lambda)\hat{j} + (5-3\lambda)\hat{k}$

Q5

Q5. $a * b = ab + 1$

i) For all $(a, b) \in (R \times R)$

$\Rightarrow ab+1 \in R$ [∵ If $a, b \in R \Rightarrow ab \in R \Rightarrow (ab+1) \in R$] [∵ Multiplication is binary operation]

∴ $a R b$ relates to a unique element in R and hence is a binary operation from $R \times R$ to R

ii) For binary operation to be associative $(a * b) * c = a * (b * c)$

LHS = $(a * b) * c$

$= (ab+1) * c$

$= (ab+1)c + 1$

$= abc + c + 1$

RHS = $a * (b * c)$

$= a * (bc+1)$

$= a(bc+1) + 1$

$= abc + a + 1$

LHS ≠ RHS

∴ It is NOT ASSOCIATIVE

Q6.

$$A = \begin{bmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{bmatrix}$$

$$A^2 = AA$$

$$= \begin{bmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{bmatrix} \begin{bmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & -1 & 2 \\ 9 & -2 & 5 \\ 0 & -1 & -2 \end{bmatrix}$$

Q7.

$$A^2 - 5A = A^2 + (-5)A$$

$$= \begin{bmatrix} 5 & -1 & 2 \\ 9 & -2 & 5 \\ 0 & -1 & -2 \end{bmatrix} + (-5) \begin{bmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & -1 & 2 \\ 9 & -2 & 5 \\ 0 & -1 & -2 \end{bmatrix} + \begin{bmatrix} -10 & 0 & -5 \\ -10 & -5 & -15 \\ -5 & 5 & 0 \end{bmatrix}$$

$$A^2 - 5A = A^2 + (-5A)$$

$$= \begin{bmatrix} +5 & -7 & -3 \\ -1 & -7 & -10 \\ -5 & 4 & -2 \end{bmatrix}$$

Q7. $I = \int \underset{I}{1} \cdot \underset{II}{\sin^{-1}(2x)}\, dx$

$$I = \left[\sin^{-1}(2x)\right](x) - \int \frac{(x)(2)}{\sqrt{1-4x^2}}\, dx \qquad \text{[INTEGRATION by parts]}$$

$$I = x\sin^{-1}(2x) - \frac{1}{2}\int \frac{2x}{\sqrt{(\frac{1}{2})^2 - x^2}}\, dx$$

$$I = x\sin^{-1}(2x) + \frac{1}{4}\int \frac{(-8x)}{\sqrt{1-4x^2}}\, dx = x\sin^{-1}(2x) + \frac{1}{4} I_1 \qquad \text{where } I_1 = \int \frac{-8x}{\sqrt{1-4x^2}}\, dx$$

Now, $I_1 = \int \frac{dt}{\sqrt{t}}$ let $1-4x^2 = t$

$-8x\,dx = dt$

$= 2\sqrt{t} + C$

$= 2\sqrt{1-4x^2} + C$

$$I = x\sin^{-1}(2x) + \frac{1}{4}\left(2\sqrt{1-4x^2}\right) + C$$

$$\boxed{I = x\sin^{-1}(2x) + \frac{\sqrt{1-4x^2}}{2} + C}$$

Q8

$$y = e^{2x}(a+bx) \quad \text{—— (1)}$$

Diff. both sides w.r.t x.

$$\frac{dy}{dx}\frac{dy}{} = e^{2x}(b) + (a+bx)(e^{2x})(2)$$

$$\Rightarrow \frac{dy}{dx}dx = e^{2x}(b+2a+2bx) \quad \text{—— (2)}$$

Diff. both sides w.r.t. x.

$$\frac{d^2y}{dx^2}\frac{d^2y}{dx^2} = e^{2x}(2b) + (b+2a+2bx)(e^{2x})(2)$$

$$= e^{2x}(4b+4a+4bx)$$

$$\frac{d^2y}{dx^2} = 2e^{2x}(2b+2a+2bx) \quad \text{—— (3)}$$

Subtract $2 \times$ eqn (2) from eqn (3)

$$\frac{d^2y}{dx^2} - 2\frac{dy}{dx} = e^{2x}(4b+4a+4bx) - e^{2x}(\text{illegible})$$

$$= e^{2x}(2b)$$

$$\Rightarrow \frac{d^2y}{dx^2} - 2\frac{dy}{dx} = 2(e^{2x})(b) \quad \text{---} (4)$$

In eq^n (3)

$$\frac{d^2y}{dx^2} = e^{2x}(4b + 4a + 4bx)$$

$$= 2\left(\frac{d^2y}{dx^2} - \frac{2dy}{dx}\right) + 4e^{2x}(a+bx) \qquad [\text{From (4)}]$$

$$\frac{d^2y}{dx^2} = \frac{2d^2y}{dx^2} - \frac{4dy}{dx} + 4y \qquad [\text{From (1)}]$$

∴ The required differential eq^n is

$$\boxed{\frac{d^2y}{dx^2} - \frac{4dy}{dx} + 4y = 0} \Rightarrow y'' - 4y' + 4y = 0.$$

Where $y'' = \frac{d^2y}{dx^2}$, $y' = \frac{dy}{dx}$, ~~$y = e^x$~~

P.T.O

Q9 $\sum_{i=0}^{2} P(X_i) = 1$ $\left[\because \sum_{i=0}^{n} P(X_i) = 1\right]$

[Sum of all probabilities = 1]

disjoint exhaustive & exclusive

$\Rightarrow k + 2k + 3k = 1$

$\Rightarrow 6k = 1$

$\boxed{k = \frac{1}{6}}$

Q11

Q10 $P(A) = \frac{3}{6} = \frac{1}{2}$

$\left[\because P(E) = \frac{\text{No. of favourable outcomes}}{\text{Total no. of possible outcomes}}\right]$

$P(B) = \frac{3}{6} = \frac{1}{2}$

$\left[\Rightarrow \text{sample space } S = \{1r, 2r, 3r, 4g, 5g, 6g\}\right.$
$A = \{2r, 4g, 6g\} \quad A \cap B = \{2r\}$
$\left.B = \{1r, 2r, 3r\}\right]$

$P(A \cap B) = P(\text{no. which is even and red}) = \frac{1}{6}$ $\left[\because \text{No. which is even and red} = \{2r\}\right]$

$P(A) \cdot P(B) = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$

Q12

$P(A \cap B) \neq P(A) \cdot P(B)$

$\therefore$ Events are not NOT INDEPENDENT.

Q11 $\vec{a} = 2\hat{i} + 3\hat{j} + \hat{k}$, $\vec{b} = \hat{i} - 2\hat{j} + \hat{k}$, $\vec{c} = -3\hat{i} + \hat{j} + 2\hat{k}$

$$[\vec{a}\ \vec{b}\ \vec{c}] = \vec{a}.(\vec{b} \times \vec{c})$$

$$= \begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{bmatrix}$$

where $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k} = 2\hat{i} + 3\hat{j} + \hat{k}$

$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k} = \hat{i} - 2\hat{j} + \hat{k}$

$\vec{c} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k} = -3\hat{i} + \hat{j} + 2\hat{k}$

$$= \begin{bmatrix} 2 & 3 & 1 \\ 1 & -2 & 1 \\ -3 & 1 & 2 \end{bmatrix}$$

$$= 2(-4-1) - 3(2+3) + 1(1-6)$$

$$= -10 - 15 - 5$$

$$\therefore \boxed{[\vec{a}\ \vec{b}\ \vec{c}] = -30}$$

Q12 $I = \int \frac{\tan^2 x \cdot \sec^2 x}{1 - \tan^6 x}\, dx$

let $\tan^3 x = t$

$3\tan^2 x \sec^2 x\, dx = dt$

Q13.

$$I = \int \frac{t^2\,dt}{1+t^6} = \int \frac{t^2}{1-(t^3)^2}\,dt$$

~~$tet \quad t^3 = m$~~

~~$3t^2 dt$~~

$$I = \frac{1}{3}\int \frac{dt}{(1)^2 - t^2}$$

$$= \frac{1}{3}\cdot\frac{1}{2.(1)}\log\left|\frac{1+t}{1-t}\right| + C \qquad \left[\because \int \frac{du}{a^2-x^2} = \log\left|\frac{a+x}{a-x}\right| + c\right]$$

$$\boxed{I = \frac{1}{6}\log\left|\frac{1+\tan^3 x}{1-\tan^3 x}\right| + c}$$

SECTION-C

P.T.O

Q13. $\tan^{-1}(2x) + \tan^{-1}(3x) = \frac{\pi}{4}$

$\tan^{-1}A + \tan^{-1}B = \tan^{-1}\left(\frac{A+B}{1-AB}\right)$, $AB < 1$

$\Rightarrow \tan^{-1}\left(\frac{5x}{1-6x^2}\right) = \frac{\pi}{4}$

taking tan on both side

$\Rightarrow \frac{5x}{1-6x^2} = 1$

$\Rightarrow 6x^2 + 5x - 1 = 0$

$\Rightarrow 6x^2 + 6x - x - 1 = 0$

$\Rightarrow (6x-1)(x+1) = 0$

$\Rightarrow x = \frac{1}{6}$ or $x = -1$

and $(2x)(3x) < 1$

$6x^2 < 1$

$\frac{-1}{\sqrt{6}} < x < \frac{1}{\sqrt{6}}$

$\therefore x = -1$ is not a solution $\left[\because -1 \notin \left(\frac{-1}{\sqrt{6}}, \frac{1}{\sqrt{6}}\right)\right]$

$\Rightarrow \boxed{x = \frac{1}{6}}$ is soln.

VERIFICATION

For $x = \frac{1}{6}$

LHS: $\tan^{-1}\left(\frac{1}{3}\right) + \tan^{-1}\left(\frac{1}{2}\right) = \tan^{-1}\left(\frac{5}{5}\right) = \frac{\pi}{4} = RHS$.

LHS = RHS

$\Rightarrow x = \frac{1}{6}$ is a soln

For $(x) = -1$

$$\tan^{-1}(-2) - \tan^{-1}(-3) = \frac{\pi}{2} - \cot^{-1}(-2) - \frac{\pi}{2} + \cot^{-1}(-3)$$

$$= \pi - \cot^{-1}(3) - \pi + \cot^{-1}(2)$$

$$= \tan^{-1}\left(\frac{1}{2}\right) - \tan^{-1}\left(\frac{1}{3}\right) = \tan^{-1}\left(\frac{\frac{1}{2}-\frac{1}{3}}{1+\frac{1}{6}}\right) = \tan^{-1}\left(\frac{1}{7}\right) \neq \text{RHS}$$

LHS $\neq$ RHS

$\therefore x = -1$ is NOT sol^n

Q15

Q14 $\log(x^2+y^2) = 2\tan^{-1}\frac{y}{x}$.

Diff. both sides w.r.t x.

$$\frac{1}{x^2+y^2}\left(2x + 2y\frac{dy}{dx}\right) = \frac{2}{1+\frac{y^2}{x^2}} \cdot \frac{d}{dx}\left(\frac{y}{x}\right)$$

$\left[\because \frac{d}{dx}(\tan^{-1}x) = \frac{1}{1+x^2}\ ;\ \frac{d}{dx}(\log x) = \frac{1}{x}\right]$

$$\Rightarrow \quad \frac{x + yy'}{x^2+y^2} = \frac{x^2}{x^2+y^2}\left(\frac{xy' - y}{x^2}\right)$$

[where $y' = \frac{dy}{dx}$]

$$\Rightarrow \quad x + yy' = xy' - y$$

$$\Rightarrow \quad x + y = y'(x - y)$$

$\Rightarrow$ $y' = \frac{x+y}{x-y}$

$$\boxed{\frac{dy}{dx} = \frac{x+y}{x-y}}$$

Hence proved.

Q15 $I = \int \frac{3x+5}{x^2+3x-18}\, dx$

$3x+5 = a\left[\frac{d}{dx}(x^2+3x-18)\right] + b$

$= a(2x+3) + b$

comparing coefficients on both sides,

$3 = 2a$ $\qquad$ $3a+b = 5$

$\boxed{a = \frac{3}{2}}$ $\qquad$ $b = 5 - 3\left(\frac{3}{2}\right) = \frac{1}{2}$ $\Rightarrow \boxed{b = \frac{1}{2}}$

P.T.O

$$I = \int \frac{\frac{3}{2}(2x+3) + \frac{1}{2}}{x^2+3x-18}\,dx$$

$$= \frac{3}{2}\int \frac{(2x+3)}{x^2+3x-18}\,dx + \frac{1}{2}\int \frac{dx}{x^2+3x-18}$$

$$= \frac{3}{2} I_1 + \frac{1}{2} I_2 .$$

$$I_1 = \int \frac{2x+3}{x^2+3x-18}\,dx \qquad I_2 = \int$$

Let $x^2+3x-18 = t$

$(2x+3)\,dx = dt$

$$I_1 = \int \frac{dt}{t} = \log|t| + c_1$$

$$I_1 = \log|x^2+3x-18| + c_1$$

Ans: $$I_2 = \int \frac{dx}{x^2+3x-18} = \int \frac{dx}{x^2+3x+\left(\frac{3}{2}\right)^2 - 18 - \left(\frac{3}{2}\right)^2}$$

$$= \int \frac{dx}{\left(x+\frac{3}{2}\right)^2 - 18 - \frac{9}{4}} = \int \frac{dx}{\left(x+\frac{3}{2}\right)^2 - \frac{81}{4}}$$

$$I_1 = -\int_a^0 f(a-t)\,dt \quad \Rightarrow$$

$$= \int_0^a f(a-t)\,dt \qquad \left[\because \int_a^b f(x)\,dx = -\int_b^a f(x)\,dx\right]$$

$$I_1 = \int_0^a f(a-x)\,dx \qquad \left[\because \int_a^b f(t)\,dt = \int_a^b f(x)\,dx\right] \quad \Rightarrow$$

LHS = RHS

Hence proved.

$$I = \int_0^{\pi} \frac{x \sin x}{1+\cos^2 x}\,dx \qquad \text{—(1)}$$

$$I = \int_0^{\pi} \frac{(\pi-x)\sin(\pi-x)}{1+\cos^2(\pi-x)}\,dx \qquad \left[\because \int_0^a f(x)\,dx = \int_0^a f(a-x)\,dx\right] \quad \therefore$$

$$I = \int_0^{\pi} \frac{(\pi-x)\sin x}{1+\cos^2 x}\,dx \qquad \text{—(2)}$$

Adding eqn (1), (2)

$$2I = \int_0^{\pi} \frac{\pi \sin x}{1+\cos^2 x}\,dx$$

$$I_2 = \int \frac{dx}{(x+\frac{3}{2})^2 - (\frac{9}{2})^2}$$

$$= \frac{1}{2 \cdot \frac{9}{2}} \log \left| \frac{(x+\frac{3}{2}) - \frac{9}{2}}{(x+\frac{3}{2}) + \frac{9}{2}} \right| + c_2$$

$$= \frac{1}{9} \log \left| \frac{x-3}{x+6} \right| + c_2$$

$$I = \frac{3}{2} \log |x^2+3x-18| + c_1 + \frac{1}{18} \log \left| \frac{x-3}{x+6} \right| + c_2$$

$$I = \frac{3}{2} \log |x^2+3x-18| + \frac{1}{18} \log \left| \frac{x-3}{x+6} \right| + c \quad \text{where } c = c_1 + c_2 = \text{constt.}$$

Q16

T.P: $\int_0^a f(x)\,dx = \int_0^a f(a-x)\,dx$

LHS: $I_1 = \int_0^a f(x)\,dx$

Let $x = a - t$

$dx = -dt$

When $x = 0 \quad t = a$

$x = a \quad t = 0$

$$\Rightarrow \quad 2I = \pi \int_0^{\pi} \frac{\sin x \, dx}{1+\cos^2 x}$$

let $\cos x = t$ — when $x = 0$ $t = 1$

$-\sin x \, dx = dt$ — $x = \pi$ $t = -1$

$$\Rightarrow \quad I = \frac{\pi}{2} \int_1^{-1} \frac{-dt}{1+t^2}$$

$$= -\frac{\pi}{2} \left[\tan^{-1} t\right]_1^{-1} = -\frac{\pi}{2} \left[-\frac{\pi}{4} - \frac{\pi}{4}\right] = \frac{\pi}{2} \cdot \frac{\pi}{2}$$

$$\boxed{I = \frac{\pi^2}{4}}$$

$$\therefore \quad \int_0^{\pi} \frac{x \sin x}{1+\cos^2 x} = \frac{\pi^2}{4}$$

Q17

$\vec{A} = \hat{i} + \hat{j} + \hat{k}$

$\vec{B} = 2\hat{i} + 5\hat{j} + 0\hat{k}$

$\vec{C} = 3\hat{i} + 2\hat{j} - 3\hat{k}$

$\vec{D} = \hat{i} - 6\hat{j} - \hat{k}$

Q.18.

$\vec{AB}$ = Position vector of $\vec{B}$ − Position vector of $\vec{A}$

$\vec{m} = \hat{i} + 4\hat{j} - \hat{k}$

$\vec{CD}$ = Position vector of $\vec{D}$ − Position vector of $\vec{C}$

$\vec{n} = -2\hat{i} - 8\hat{j} + 2\hat{k}$

Angle b/w $\vec{AB}$ and $\vec{CD}$ $= \theta$

$$\cos\theta = \frac{\vec{m}.\vec{n}}{|\vec{m}||\vec{n}|}$$

$$= \frac{m_1n_1 + m_2n_2 + m_3n_3}{\sqrt{m_1^2+m_2^2+m_3^2}\sqrt{n_1^2+n_2^2+n_3^2}}$$

$$= \frac{-2-32-2}{\sqrt{1+16+1}\sqrt{4+64+4}}$$

$$= \frac{-36}{\sqrt{18}\sqrt{72}} = \frac{-36}{(3\sqrt{2})(6\sqrt{2})} = \frac{-36}{18\times 2}$$

$$= \frac{-36}{36} = -1$$

$\theta = \cos^{-1}(-1) \Rightarrow \boxed{\theta = \pi}$

$\therefore$ Since, $\vec{AB}$ and $\vec{CD}$ are antiparallel,

they ARE COLLINEAR.

Q18.

$$\Delta = \begin{vmatrix} a+b+c & -c & -b \\ -c & a+b+c & -a \\ -b & -a & a+b+c \end{vmatrix}$$

$$\Delta = \begin{vmatrix} a+b & a+b & -(a+b) \\ -c & a+b+c & -a \\ -b & -a & a+b+c \end{vmatrix}$$ $R_1 \to R_1 + R_2$

$$= (a+b) \begin{vmatrix} 1 & 1 & -1 \\ -c & a+b+c & -a \\ -b & -a & a+b+c \end{vmatrix}$$ Taking $(a+b)$ common from R_1

$$= (a+b) \begin{vmatrix} 1 & 1 & -1 \\ -(b+c) & b+c & b+c \\ -b & -a & a+b+c \end{vmatrix}$$ $R_2 \to R_2 + R_3$

$$= (a+b)(b+c) \begin{vmatrix} 1 & 1 & -1 \\ -1 & 1 & 1 \\ -b & -a & a+b+c \end{vmatrix}$$ Taking $(b+c)$ common from R_2

$$= (a+b)(b+c) \begin{vmatrix} 0 & 0 & -1 \\ 0 & 2 & 1 \\ a+c & b+c & a+b+c \end{vmatrix}$$ $C_1 \to C_1 + C_3$, $C_2 \to C_2 + C_3$

$\Delta = (a+b)(b+c)(c+a)$ $\begin{vmatrix} 0 & 0 & -1 \\ 0 & 2 & 1 \\ 1 & b+c & a+b+c \end{vmatrix}$ Taking $(a+c)$ common from C_1

Expanding along C_1

$\Delta = (a+b)(b+c)(c+a)(2)$

$\Delta = 2(a+b)(b+c)(c+a)$

Hence proved.

Q19 $y = \sin t$ $x = \cos t + \log(\tan(t/2))$

Diff. wr.t t

$\frac{dy}{dt} = \cos t$ ——(1)

Diff. wr.t t

$\frac{dx}{dt} = -\sin t + \frac{1}{\tan t/2} \cdot \sec^2 t/2 \cdot \frac{1}{2}$

$= -\sin t + \frac{\cos t/2}{2 \sin t/2 \cos^2 t/2}$

$= -\sin t + \frac{1}{\sin t}$ [$\because 2\sin\theta\cos A = \sin 2A$]

Diff w.r.t dt

$\frac{d^2y}{dt^2} = -\sin t$

$\left.\frac{d^2y}{dt^2}\right]_{t=\pi/4} = -\sin\frac{\pi}{4}$


$\left.\frac{d^2y}{dt^2}\right]_{t=\pi/4} = -\frac{1}{\sqrt{2}}$


$\frac{dx}{dt} = -\sin t + \frac{1}{\sin t}$ ——(2)

$\frac{dx}{dt} = \frac{\cos^2 t}{\sin t}$

P.T.O

Q 29


SCALE
On x axis: 1 small = 1 unit of model A
On y axis: 1 small = 1 unit of model B
= Feasible region
No of model B items
No of model A items
Y
Y'
X
X'
(0, 40)
2x+y≤40
x≥0
(0, 26.7)
(10, 20)
2x+3y≤80
(20, 0)
(40, 0)
y≥0
0
10
20
30
40
50
60
70
80
90
100
110



Q19 Contd...

~~Diff~~ Divide eqn (1) by (2)

$$\frac{dy/dt}{dx/dt} = \frac{\cos t}{-\sin t + \frac{1}{\sin t}}$$

$$\frac{dy}{dx} = \frac{\cos t \cdot \sin t}{1-\sin^2 t} = \frac{\sin t}{\cos t} = \tan t$$

Diff. w.r.t x.

$$\frac{d^2y}{dx^2} = \frac{d(\tan t)}{dt} \frac{dt}{dx}$$

$$= \sec^2 t \cdot \frac{1}{dx/dt} = \frac{\sec^2 t}{-\sin t + \frac{1}{\sin t}}$$

$$= \frac{\sec^2 t \cdot \sin t}{1-\sin^2 t} = \frac{\sin t}{\cos^2 t \cdot \cos^2 t} = \frac{\sin t}{\cos^4 t}$$

$$= \sin t \cdot \sec^4 t$$

$$\left.\frac{d^2y}{dx^2}\right|_{t=\pi/4} = \frac{1}{\sqrt{2}} (\sqrt{2})^4 = \frac{4}{\sqrt{2}} = 2\sqrt{2}$$

$$\therefore \left.\frac{d^2y}{dx^2}\right|_{t=\pi/4} = 2\sqrt{2}$$

Q20.

$R = \{(a,b) : a \leq b\}$

REFLEXIVE:

Every element $a \in \mathbb{R}$ is equal to itself

$\Rightarrow a = a$

$\Rightarrow a \leq a$ is true

$\therefore (a,a) \in R$ for all $a \in \mathbb{R}$

The relation is REFLEXIVE

where $\mathbb{R}$ = set of all Real nos

R = Relation

TRANSITVE:

For all $(a,b) \in R$ and $(b,c) \in R$

$a \leq b$ and $b \leq c$

$\Rightarrow a \leq c$

$\Rightarrow (a,c) \in R$

where $a, b, c \in \mathbb{R}$

$\therefore$ The ~~set~~ is tr relation is TRANSITVE

$\because$ for $(a,b), (b,c) \in R$, ~~$(a,c) \in R$~~ $(a,c) \in R$

Q21.

SYMMETRIC : For relation to be symmetric,
for all $(a, b) \in R$, (b, a) should also exist in R

~~$a \leq b \not\Rightarrow$~~ $a \leq b$

$b \not\leq a$ → This relation is true only $a = b = 1$.

For eg: $\frac{1}{2} \leq 1 \Rightarrow (\frac{1}{2}, 1) \in R$

but $1 \not\leq \frac{1}{2}$ $\therefore (1, \frac{1}{2}) \notin R$

$\therefore$ Relation is NOT SYMMETRIC

Q21 $y = \sqrt{3x-2}$

$\left.\frac{dy}{dx}\right|_{(x_1,y_1)} = \frac{3}{2\sqrt{3x_1-2}} = \frac{3}{2y_1}$ = slope of tangent = m

Acc. to question $= 2$

$\Rightarrow \frac{3}{2y_1} = 2 \Rightarrow \boxed{y_1 = \frac{3}{4}}$

[$\because$ slope of $(4x-2y+5=0)$ is $m_1 = 2$. and ||el line have equal slope]

$\Rightarrow \frac{3}{2\sqrt{3x_1-2}} = 2 \Rightarrow 3 = 4\sqrt{3x_1-2}$

$\Rightarrow 9 = 16(3x_1-2)$ squaring both side

$\Rightarrow 3x_1 - 2 = \frac{9}{16} \Rightarrow x_1 = \frac{41}{48} \Rightarrow \boxed{x_1 = \frac{41}{48}}$

$x_1 = \frac{41}{48}$

$\Rightarrow y_1 = \sqrt{3x_1 - 2} = \sqrt{3 \times \frac{41}{48} - 2} = \sqrt{\frac{41 - 32}{16}} = \pm\frac{3}{4} = \left|\frac{3}{4}\right| = \sqrt{\frac{9}{16}}$

At $\left(\frac{41}{48}, \frac{3}{4}\right)$ and $\left(\frac{41}{48}, \frac{-3}{4}\right)$, ~~slope of~~ tangent is parallel to given line.

(But $\because y \not< 0 \Rightarrow y \geq 0$) ↑ Not possible

EQUATION OF TANGENT

For $x_1 = 41/48$, $y_1 = 3/4$

$$y - \frac{3}{4} = 2\left(x - \frac{41}{48}\right)$$

$$y - \frac{3}{4} = 2x - \frac{41}{24}$$

$$2x - y - \frac{41}{24} + \frac{18}{24} = 0$$

$$\Rightarrow 2x - y - \frac{23}{24} = 0$$

$$48x - 24y - 23 = 0$$

For $x_1 = 41/48$, $y_1 = 3/4$

$$y + \frac{3}{4} = 2\left(x - \frac{41}{48}\right)$$

$$2x - y - \frac{41}{24} - \frac{18}{24} = 0$$

$$48x - 24y - 59 = 0$$

↑ NOT POSSIBLE

X

Slope of normal $= \frac{-1}{\left[\frac{dy}{dx}\right]_{(x_1,y_1)}} = \frac{-1}{2}$

EQUATION OF NORMAL :

At $\left(\frac{41}{48}, \frac{3}{4}\right)$

$$y - \frac{3}{4} = -\frac{1}{2}\left(x - \frac{41}{48}\right)$$

$$\Rightarrow y - \frac{3}{4} = -\frac{x}{2} + \frac{41}{96}$$

$$\Rightarrow y + \frac{x}{2} - \frac{3}{4} - \frac{41}{96} = 0$$

$$\Rightarrow y + \frac{x}{2} + \left(\frac{-72-41}{96}\right) = 0$$

$$\Rightarrow \boxed{96y + 48x - 113 = 0}$$

At $\left(\frac{41}{48}, -\frac{3}{4}\right)$

$$y + \frac{3}{4} = -\frac{1}{2}\left(x - \frac{41}{48}\right)$$

$$\Rightarrow y + \frac{3}{4} + \frac{x}{2} - \frac{41}{96} = 0$$

$$\Rightarrow y + \frac{x}{2} + \frac{72-41}{96} = 0$$

$$\Rightarrow \boxed{96y + 48x + 31 = 0}$$

NOT POSSIBLE

P.T.O.

Q22. $(1+x^2)\frac{dy}{dx} + 2xy = 4x^2$

$$\frac{dy}{dx} + \left(\frac{2x}{1+x^2}\right)y = \frac{4x^2}{1+x^2}$$

It is linear DE of form $\frac{dy}{dx} + Py = Q$

$P = \frac{2x}{1+x^2}$ $\quad Q = \frac{4x^2}{1+x^2}$

Q23.

$$I.F = e^{\int P\,dx} = e^{\ln|1+x^2|} = 1+x^2$$

Soln of DE:

$$y(1+x^2) = \int \frac{4x^2}{1+x^2}(1+x^2)\,dx + C$$

$$= \int 4x^2\,dx + C$$

$$\boxed{y(1+x^2) = \frac{4x^3}{3} + C}$$

$x=0, y=0$

$\therefore c=0$

$\Rightarrow \boxed{3y(1+x^2)=4x^3}$

Q23

Line I

$$\frac{1-x}{3}=\frac{7y-14}{\lambda}=\frac{z-3}{2}$$

$$\Rightarrow \frac{x-1}{-3}=\frac{y-2}{\lambda/7}=\frac{z-3}{2}=\beta$$

Line II

$$\frac{7-7x}{3\lambda}=\frac{y-5}{1}=\frac{6-z}{5}$$

$$\frac{x-1}{-3\lambda/7}=\frac{y-5}{1}=\frac{z-6}{-5}=\alpha$$

Direction ratios are $\left(-3, \frac{\lambda}{7}, 2\right)$ (a_1, a_2, a_3) and $\left(\frac{-3\lambda}{7}, 1, -5\right)$ (b_1, b_2, b_3) respectively

For lines to be $\perp$, ~~$\vec{a_1}\cdot\vec{a_2}=0$~~ $\vec{a_1}\cdot\vec{b}=0$ where ($\vec{a}, \vec{b}$ are ||el vectors of lines)

$$a_1b_1+a_2b_2+a_3b_3=0$$

$$\frac{9\lambda}{7}+\frac{\lambda}{7}-10=0$$

$$\frac{10\lambda}{7}=10$$

$$\boxed{\lambda=7}$$

$\therefore$ For $\lambda=7$, lines are $\perp$.

Q24.

Any point of line I is $(-3\beta+1,\ \beta+2,\ 2\beta+3)$

Line II is $(-3\mu+1,\ \mu+5,\ -5\mu+6)$

For lines to intersect, they should be equal.

$-3\beta+1 = -3\mu+1$ $\quad$ $\mu+5=\beta+2$ $\quad$ $2\beta+3 = -5\mu+6$

$3\mu-3\beta=0$ $\quad$ $\mu-\beta+3=0$ — (3) $\quad$ $2\beta+5\mu=3$ — (2)

$\mu=\beta$ — (1)

From (1), (2)

$7\mu=3$

$\boxed{\mu=\frac{3}{7}}$ $\quad$ $\boxed{\beta=\frac{3}{7}}$

In eqn (3)

$\frac{3}{7}-\frac{3}{7}+3 \neq 0$


$\therefore$ Values do not satisfy eqn (3)

$\Rightarrow$ LINES <u>do NOT</u> intersect.

P.T.O

SECTION-D

Q24.



Consider a sphere of radius r.

Then, height of cone = $r + x = H$

Radius of cone = $\sqrt{r^2 - x^2} = R$

Volume of cone = $V = \frac{1}{3}\pi R^2 H$

$V = \frac{1}{3}\pi (r^2 - x^2)(r+x)$

$= \frac{\pi}{3}(r^3 - rx^2 + xr^2 - x^3)$

$\frac{dV}{dx} = \frac{\pi}{3}(-2rx + r^2 - 3x^2)$

For critical pt $\frac{dV}{dx} = 0$

$r^2 - 2rx - 3x^2 = 0$

$r^2 - 3rx + rx - 3x^2 = 0$

$(r - x)(r - 3x) = 0$

x cannot be -ve $\Rightarrow \boxed{r = x}$ is a critical pt

$\boxed{\frac{r}{3}}$

$$\frac{d^2V}{dx^2} = \frac{\pi}{3}(-2r-6x)$$

$$\left.\frac{d^2V}{dx^2}\right]_{x=r/3} = \frac{\pi}{3}[-2r-2r] = \frac{-4r\pi}{3} < 0$$

Using Second Derivative test, $\therefore \boxed{x = \frac{r}{3}}$ is a point of maxima

$H = r + x$

$$\boxed{H = \frac{4r}{3}}$$ Hence proved

Max volume of cone $= V_{max} = \frac{\pi}{3}(r+x)(r^2-x^2)$

$$= \frac{\pi}{3}\left(\frac{4r}{3}\right)\left(r^2 - \frac{r^2}{9}\right)$$

$$= \frac{4\pi r^3 \cdot 8}{81} = \frac{32\pi r^3}{27} \text{ (unit)}^3$$

$$\boxed{V_{max} = \frac{32\pi r^3}{81} \text{ (unit)}^3}$$

Q25.

$$A = \begin{bmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{bmatrix}$$

$$\text{adj } A = \begin{bmatrix} 0 & +2 & 1 \\ -1 & -9 & -5 \\ 2 & +23 & 13 \end{bmatrix}^T = \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix}$$

$$|A| = 2(0) + 3(-2) + 5(1)$$

$$= 0 - 6 + 5 = -1$$

$$A^{-1} = \frac{1}{|A|} \text{adj } A = -1 \begin{bmatrix} 0 & -1 & 2 \\ 2 & -9 & 23 \\ 1 & -5 & 13 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} 0 & 1 & -2 \\ -2 & 9 & -23 \\ -1 & 5 & -13 \end{bmatrix}$$

Given Eqⁿ:

$$2x - 3y + 5z = 11$$

$$3x + 2y - 4z = -5$$

$$x - y - 2z = -3$$

$AX = B$

Pre Premultiply A^{-1} on both sides

$A^{-1}AX = A^{-1}B$

$IX = A^{-1}B$

$X = A^{-1}B$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 & 1 & -2 \\ -2 & 9 & -23 \\ -1 & 5 & -13 \end{bmatrix} \begin{bmatrix} 11 \\ -5 \\ -3 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -5+6 \\ -22-45+69 \\ -11-25+39 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$x = 1$

$y = 2$

$z = 3$

026

026

Q26.

A = Event of choosing a defective item.

E_1 = Item produced by A

E_2 = Item produced by B

E_3 = Item produced by C.

$P(E_1) = \frac{50}{100} = \frac{1}{2}$, $P(E_2) = \frac{30}{100} = \frac{3}{10}$, $P(E_3) = \frac{20}{100} = \frac{1}{5}$

$P\left(\frac{A}{E_1}\right) = \frac{1}{100}$, $P\left(\frac{A}{E_2}\right) = \frac{5}{100}$; $P\left(\frac{A}{E_3}\right) = \frac{7}{100}$

Using Baye's Thm.,

$$A\left(\frac{E_1}{A}\right) = \frac{P(E_1)\,P(A/E_1)}{P(E_1)\,P(A/E_1) + P(E_2)\,P\left(\frac{A}{E_2}\right) + P(E_3)\cdot P(A/E_3)}$$

$$= \frac{\frac{1}{2}\times\frac{1}{100}}{\frac{1}{2}\times\frac{1}{100} + \frac{3}{10}\times\frac{5}{100} + \frac{1}{5}\times\frac{7}{100}}$$

$$= \frac{\frac{1}{200}}{\frac{1}{200} + \frac{3}{200} + \frac{7}{500}} = \frac{\frac{1}{2}}{\frac{1}{2} + \frac{3}{2} + \frac{7}{5}} = \frac{\frac{1}{2}}{2 + \frac{7}{5}} = \frac{\frac{1}{2}}{\frac{17}{5}}$$

$P(E_1/A) = \frac{5}{34}$ = Probability that defective item was produced by A

Q27. Let $\vec{n} = A\hat{i} + B\hat{j} + C\hat{k}$ of the required plane

Eqn of plane is $A(x-2) + B(y-2) + C(z+1) = 0$ — (3)

The plane also passes through (3,4,2) & (7,0,6)

$$A + 2B + 3C = 0 \quad — (1)$$

$$5A - 2B + 7C = 0 \quad — (2)$$

Adding (1), (2)

$$6A + 10C = 0$$

$$\boxed{A = -\frac{5C}{3}}$$

In eqn(2) $\frac{-25C}{3} + 7C = 2B$

$$\frac{-4C}{2\times 3} = B$$

$$\Rightarrow \boxed{B = -\frac{2C}{3}}$$

$$\vec{n_1} = -\frac{5C}{3}\hat{i} - \frac{2C}{3}\hat{j} + C\hat{k} = -\frac{C}{3}\left(5\hat{i} + 2\hat{j} - 3\hat{k}\right)$$

Q29. Let x be no. of A models
y be no. of B models.

Objective $Z = 15x + 10y$ (Maximize)

Subject to constraints:

$x, y \geq 0$

$2x + y \leq 40$ (skilled man working hrs)

$2x + 3y \leq 80$ (Semi-skilled man working hrs)

$2x + y \leq 40$

$2x + y = 40$

x	0	20
y	40	0

Zero test : TRUE

$2x + 3y \leq 80$

$2x + 3y = 80$

x	0	40
y	80/3	0

Zero test : TRUE

GRAPH: On graph paper.

Eqⁿ of line BC:

$(y-2) = \frac{5}{-2}(x-6)$

$\Rightarrow y-2 = -\frac{5}{2}x + 15$

$\Rightarrow \boxed{y_2 = -\frac{5}{2}x + 17}$

Eqⁿ of line AC

$(y-5) = \frac{3}{-4}(x-2)$

$y-5 = -\frac{3}{4}x + \frac{3}{2}$

$\boxed{y_3 = -\frac{3x}{4} + \frac{13}{2}}$

Area of shaded region is required area A = ar(ABDE) + ar(BCFD) − ar(ACFE)

$$A = \int_2^4 y_1\,dx + \int_4^6 y_2\,dx - \int_2^6 y_3\,dx$$

$$= \int_2^4 (x+3)\,dx + \int_4^6 \left(\frac{-5x}{2}+17\right)dx - \int_2^6 \left(\frac{-3x}{4}+\frac{13}{2}\right)dx$$

$$A = \left[\frac{x^2}{2}+3x\right]_2^4 + \left[17x - \frac{5x^2}{4}\right]_4^6 - \left[\frac{13x}{2} - \frac{3x^2}{8}\right]_2^6$$

$$= \left[8+12-2-6\right] + \left[102 - 45 - 68 + 20\right] - \left[39 - 27 - 13 + \frac{3}{2}\right]$$

$$= 12 + 9 - 14$$


$$\boxed{A = 7 \text{ sq. units}}$$

∴ Eqⁿ of other plane is

$\vec{r}.(5\hat{i}+2\hat{j}-3\hat{k}) = 23$ → vector eqⁿ ⇒ $\vec{r}.(5\hat{i}+2\hat{j}-3\hat{k}) - 23 = 0$

$5x+2y-3z = 23$ → cartesian eqⁿ

Q28.



Eqⁿ of general line:

$$y-y_1 = \frac{y_2-y_1}{x_2-x_1}(x-x_1)$$

Eqⁿ of line AB :

$$(y-5) = \frac{2}{2}(x-2)$$

$y = x+3$

Direction ratios of normal to the plane are

$\left(\frac{-5c}{3}, \frac{-2c}{3}, c\right) \equiv$ ~~$(-5c, -2c, 3c)$~~ $(5, 2, -3)$

$\therefore \vec{n} = 5\hat{i} + 2\hat{j} - 3\hat{k}$

Eq^n: of plane: $5(x-2) + 2(y-2) - 3(z+1) = 0$ [From (3)]

$5x + 2y - 3z - 10 - 4 - 3 = 0$

$\boxed{5x + 2y - 3z = 17}$ is cartesian eq^n of plane.

VECTOR EQN:

$\boxed{\vec{r}.\vec{n} = d} \Rightarrow \left[\vec{r} - (2\hat{i} + 2\hat{j} - \hat{k})\right].\vec{n} = 0$ [$\because (\vec{r} - \vec{a}).\vec{n} = 0$ is eq^n of plane]

$\vec{r}(5\hat{i} + 2\hat{j} - 3\hat{k}) - (2\hat{i} + 2\hat{j} - \hat{k})(5\hat{i} + 2\hat{j} - 3\hat{k}) = 0$

$\Rightarrow \boxed{\vec{r}(5\hat{i} + 2\hat{j} - 3\hat{k}) = 17}$ is vector eq^n of plane.

For plane parallel to above plane, $\vec{n_2} = \vec{n} = 5\hat{i} + 2\hat{j} - 3\hat{k}$

$(\vec{r} - \vec{a_1}).\vec{n} = 0$

Here $\vec{a_1} = 4\hat{i} + 3\hat{j} + \hat{k}$

$\left[\vec{r} - (4\hat{i} + 3\hat{j} + \hat{k})\right].(5\hat{i} + 2\hat{j} - 3\hat{k}) = 0$

$\vec{r}(5\hat{i} + 2\hat{j} - 3\hat{k}) - (20 + 6 - 3) = 0$

Corner pt	$Z = 15x + 10y$	
$(0,0)$	$Z = 0$	
~~$(\frac{80}{3}, 0)$~~ $(0, \frac{80}{3})$	$Z = 400 - 0 + \frac{800}{3} = 266.7 \simeq 267$	
$(10, 20)$	$Z = 150 + 200 = 350$	(MAX)
$(20, 0)$	$Z = 300$	

∴ No. of model A = 10

No. of model B = 20

Maximum profit = ₹ 350

